


Third Midterm: Solutions

1. Rewrite the given equation in the form $\frac{x^2}{\frac{2}{3}} + \frac{y^2}{\frac{2}{3}} + \frac{z^2}{1} = 1$. This equation defines an ellipsoid centered at the origin with the x -, y - and z - radii equal to $\sqrt{\frac{2}{3}}$, $\sqrt{\frac{2}{3}}$ and 1 respectively. It looks as follows:



Such an ellipsoid can be parametrized by


$$x(\theta, \phi) = \sqrt{\frac{2}{3}} \sin \phi \cos \theta, \quad y(\theta, \phi) = \sqrt{\frac{2}{3}} \sin \phi \sin \theta, \quad z(\theta, \phi) = \cos \phi. \quad (1)$$

It is important to notice here that the parameter ϕ is **not** the angle between a radius-vector and the z -axis.

The condition $z \geq x^2 + y^2$ puts certain restrictions on ϕ . To determine them, we find the intersection of the given ellipsoid and the surface¹ $z = x^2 + y^2$:

$$\begin{cases} 3(x^2 + y^2) + 2z^2 = 2 \\ z = x^2 + y^2. \end{cases}$$


We plug $z = x^2 + y^2$ into the first equation and obtain $2z^2 + 3z + 2 = 0$. The positive solution of this quadratic equation is $z = \frac{1}{2}$. Now, we plug $z = \frac{1}{2}$ into the first equation of the system again and obtain $x^2 + y^2 = \frac{1}{2}$. Hence, the intersection is a circle of radius $\frac{1}{\sqrt{2}}$ lying on the plane $z = \frac{1}{2}$.



¹It is a paraboloid.

If $z = \frac{1}{2}$, then, according to (1), $\phi = \cos^{-1}\left(\frac{1}{2}\right) = \frac{\pi}{3}$. Therefore, ϕ ranges from 0 to $\frac{\pi}{3}$ and θ goes from 0 to 2π .

This is how the surface looks like:



2. We denote the given vector field $6xy(\cos z)\mathbf{i} + 3x^2(\cos z)\mathbf{j} - 3x^2y(\sin z)\mathbf{k}$ by $\mathbf{F} = (F_1, F_2, F_3)$. All F_i 's are continuously differentiable, and it is easy to verify that $\nabla \times \mathbf{F} = \mathbf{0}$. Hence, \mathbf{F} is a conservative field and we can be sure that a function f satisfying $\mathbf{F} = \nabla f$ exists.

We have

$$\begin{aligned} f(x, y, z) &= \int F_1 dx = \int 6xy(\cos z) dx = 3x^2y(\cos z) + h_1(y, z), \\ f(x, y, z) &= \int F_2 dy = \int 3x^2(\cos z) dy = 3x^2y(\cos z) + h_2(x, z), \\ f(x, y, z) &= \int F_3 dz = \int -3x^2y(\sin z) dz = 3x^2y(\cos z) + h_3(x, y). \end{aligned}$$

Thus, f can be taken in the form $f(x, y, z) = 3x^2y(\cos z)$.

3. First, we find

$$\Phi_u(u, v) = (1, 1, v), \quad \Phi_v(u, v) = (-1, 1, u).$$

The cross product $\Phi_u(1, 1) \times \Phi_v(1, 1)$ gives a vector which is normal to the given surface at the point $\Phi(1, 1) = (0, 2, 1)$. We calculate

$$\begin{aligned} \Phi_u(u, v) \times \Phi_v(u, v) &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & v \\ -1 & 1 & u \end{vmatrix} = (u - v)\mathbf{i} - (u + v)\mathbf{j} + 2\mathbf{k} \\ \Phi_u(1, 1) \times \Phi_v(1, 1) &= (0, -2, 2). \end{aligned}$$

This vector can be taken as a normal vector of the tangent plane at the point $\Phi(1, 1)$. Hence, an equation of the tangent plane is

$$0(x - 0) - 2(y - 2) + 2(z - 1) = 0.$$

That finishes the first part.

$$\begin{aligned} \text{Area} &= \int \int_S 1 \, dS = \int \int_D 1 \cdot \|\Phi_u(u, v) \times \Phi_v(u, v)\| \, du \, dv \\ &= \int \int_D \sqrt{(u-v)^2 + (u+v)^2 + 2^2} \, du \, dv = \sqrt{2} \int \int_D \sqrt{u^2 + v^2 + 2} \, du \, dv, \end{aligned}$$

where D is the unit disk.

To calculate the latter integral we use the polar coordinates. We let $u = r \cos \theta$ and $v = r \sin \theta$. Then

$$\begin{aligned} \sqrt{2} \int \int_D \sqrt{u^2 + v^2 + 2} \, du \, dv &= \sqrt{2} \int_0^{2\pi} \int_0^1 \sqrt{r^2 + 2} \cdot \underbrace{r}_{\text{Jacobian}} \, dr \, d\theta = \sqrt{2} \cdot 2\pi \cdot \left. \frac{(r^2 + 2)^{\frac{3}{2}}}{3} \right|_0^1 \\ &= \frac{2\sqrt{2}\pi}{3} (3\sqrt{3} - 2\sqrt{2}). \end{aligned}$$

4. The given cylinder can be parametrized by

$$\mathbf{T}(\theta, z) = (\cos \theta, \sin \theta, z), \quad 0 \leq \theta \leq 2\pi, \quad 0 \leq z \leq 1.$$

We calculate


$$\begin{aligned} \mathbf{T}_\theta(\theta, z) &= (-\sin \theta, \cos \theta, 0) \\ \mathbf{T}_z(\theta, z) &= (0, 0, 1) \\ \mathbf{T}_\theta(\theta, z) \times \mathbf{T}_z(\theta, z) &= (\cos \theta, \sin \theta, 0). \end{aligned}$$

$$\begin{aligned} \int \int_S \mathbf{F} \, d\mathbf{S} &= \int \int_S \mathbf{F} \cdot (\mathbf{T}_\theta \times \mathbf{T}_z) \, dS = \int_0^{2\pi} \int_0^1 (\cos \theta, \sin \theta, -\sin \theta) \cdot (\cos \theta, \sin \theta, 0) \, dz \, d\theta \\ &= \int_0^{2\pi} \int_0^1 (\cos^2 \theta + \sin^2 \theta) \, dz \, d\theta = 2\pi. \end{aligned}$$

5. The change of variables formula yields

$$\int \int_B x^2 + y^2 \, dx \, dy = \int \int_{B^*} (x(u, v)^2 + y(u, v)^2) \cdot \left| \frac{\partial(x, y)}{\partial(u, v)} \right| \, du \, dv, \quad (2)$$

where $B^* = [1, 4] \times [1, 3]$.



We know exactly how u and v depend on x and y , but expressing x and y in terms of u, v seems to be a not so easy problem. It suggests to look for an alternative way of calculating the Jacobian $\left| \frac{\partial(x,y)}{\partial(u,v)} \right|$. We will use the following property of Jacobians:

$$\left| \frac{\partial(x,y)}{\partial(u,v)} \right| = \left| \frac{\partial(u,v)}{\partial(x,y)} \right|^{-1}. \quad (3)$$

We find

$$\left| \frac{\partial(u,v)}{\partial(x,y)} \right| = \begin{vmatrix} 2x & -2y \\ y & x \end{vmatrix} = 2(x^2 + y^2).$$

Then, according to the identity (3),

$$\left| \frac{\partial(x,y)}{\partial(u,v)} \right| = \frac{1}{2(x(u,v)^2 + y(u,v)^2)}$$

Returning to (2), we obtain

$$\begin{aligned} \iint_B x^2 + y^2 \, dx \, dy &= \iint_{B^*} (x(u,v)^2 + y(u,v)^2) \cdot \frac{1}{2(x(u,v)^2 + y(u,v)^2)} \, du \, dv \\ &= \frac{1}{2} \iint_{B^*} 1 \, du \, dv = \frac{1}{2} \cdot \text{Area of } B^* = 3. \end{aligned}$$


We still do not know the formulas for $x(u,v)$, $y(u,v)$, but nevertheless, we calculated the wanted integral using the change of variables.

6. Short Solution.

By Stokes' Theorem,

$$\int_{C=\partial T} \mathbf{F} \, d\mathbf{S} = \int \int_T (\nabla \times \mathbf{F}) \, d\mathbf{S},$$

where $\mathbf{F} = (x+y, 2x-z, y+z)$ and T is the triangle with the vertices $(2, 0, 0)$, $(0, 3, 0)$, $(0, 0, 6)$.



We calculate

$$\nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x + y & 2x - z & y + z \end{vmatrix} = 2\mathbf{i} + \mathbf{k}.$$

Then

$$\int \int_T (\nabla \times \mathbf{F}) \, d\mathbf{S} = \int \int_T (\nabla \times \mathbf{F}) \cdot \mathbf{n} \, dS = \int \int_T (2, 0, 1) \cdot \mathbf{n} \, dS, \quad (4)$$

where \mathbf{n} is the unit normal vector to the plane containing the triangle T pointing “upwards” (it is chosen with respect to the right-hand rule).

To find \mathbf{n} , let

$$\mathbf{a} = (0, 3, 0) - (2, 0, 0) = (-2, 3, 0)$$

$$\mathbf{b} = (0, 0, 6) - (2, 0, 0) = (-2, 0, 6)$$

and calculate

$$\mathbf{a} \times \mathbf{b} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -2 & 3 & 0 \\ -2 & 0 & 6 \end{vmatrix} = 18\mathbf{i} + 12\mathbf{j} + 6\mathbf{k}.$$

Then $\mathbf{n} = \frac{\mathbf{a} \times \mathbf{b}}{\|\mathbf{a} \times \mathbf{b}\|}$. Returning to (4), we obtain

$$\begin{aligned} \int \int_T (\nabla \times \mathbf{F}) \, d\mathbf{S} &= \int \int_T (2, 0, 1) \cdot \frac{\mathbf{a} \times \mathbf{b}}{\|\mathbf{a} \times \mathbf{b}\|} \, dS = \frac{1}{\|\mathbf{a} \times \mathbf{b}\|} \int \int_T (2, 0, 1) \cdot (18, 12, 6) \, dS \\ &= \frac{42}{\|\mathbf{a} \times \mathbf{b}\|} \cdot \int \int_T 1 \, dS = \frac{42}{\|\mathbf{a} \times \mathbf{b}\|} \cdot \text{Area of } T. \end{aligned}$$

Recall that the area of a triangle with the sides \mathbf{a} , \mathbf{b} is equal to $\frac{1}{2}\|\mathbf{a} \times \mathbf{b}\|$ and conclude that the latter integral is equal to 21.

Long Solution.

Parametrize each side of the triangle, calculate the line integral of \mathbf{F} over each side and sum up the results afterwards.