

VERSION D

MATH 1271 Fall 2011, Midterm #2
Handout date: Thursday 10 November 2011

PRINT YOUR NAME:

SOLUTIONS

PRINT YOUR TA'S NAME:

WHAT SECTION ARE YOU IN?

Closed book, closed notes, no calculators/PDAs; no reference materials of any kind. Turn off all handheld devices, including cell phones.

Show work; a correct answer, by itself, may be insufficient for credit. Arithmetic need not be simplified, unless the problem requests it.

I understand the above, and I understand that cheating has severe consequences, from a failing grade to expulsion.

SIGN YOUR NAME:

I. Multiple choice

A. (5 pts) (no partial credit) Find the logarithmic derivative of $(2 + \sin x)^x$ w.r.t. x .

(a) $\cos x$

(b) $\ln(\cos x)$

(c) $\ln(2 + \sin x) + \left(\frac{x \cos x}{2 + \sin x}\right)$

$$\frac{d}{dx} \left[x (\ln(2 + \sin x)) \right]$$

(d) $[(2 + \sin x)^x] \left[\ln(2 + \sin x) + \left(\frac{x \cos x}{2 + \sin x}\right) \right]$

(e) NONE OF THE ABOVE

B. (5 pts) (no partial credit) Find the derivative of $(2 + \sin x)^x$ w.r.t. x .

(a) $\cos x$

(b) $\ln(\cos x)$

(c) $\ln(2 + \sin x) + \left(\frac{x \cos x}{2 + \sin x}\right)$

(d) $[(2 + \sin x)^x] \left[\ln(2 + \sin x) + \left(\frac{x \cos x}{2 + \sin x}\right) \right]$

(e) NONE OF THE ABOVE

C. (5 pts) (no partial credit) Compute $\lim_{x \rightarrow 0} \left[\frac{\sin^2 x}{4x^3 + 2x^2} \right]$.

(a) 2

(b) 1

(c) $1/2$

(d) $1/4$

(e) NONE OF THE ABOVE

$$\frac{x^2}{2x^2} = \frac{1}{2} \longrightarrow \frac{1}{2}$$

D. (5 pts) (no partial credit) Find the slope of the tangent line to $y = (x^3 + 4)e^{2x}$ at the point $(0, 4)$.

- (a) 2
- (b) 4
- (c) 6
- (d) 8
- (e) NONE OF THE ABOVE

$$\begin{aligned} & \left[3x^2 \cdot e^{2x} + (x^3 + 4) \cdot e^{2x} \cdot 2 \right]_{x \rightarrow 0} \\ & = 0 + 4 \cdot e^0 \cdot 2 \\ & = 8 \end{aligned}$$


E. (5 pts) (no partial credit) Find the logarithmic derivative of $x^2 + 3x - 8$ w.r.t. x .

- (a) $\frac{2x + 3}{x^2 + 3x - 8}$
- (b) $\frac{x^2 + 3x - 8}{2x + 3}$
- (c) $(\ln(x^2)) + 3(\ln x) - (\ln 8)$
- (d) $\ln(2x + 3)$
- (e) NONE OF THE ABOVE

F. (5 pts) (no partial credit) Suppose $f'(x) = -x^2 + 3x - 2$. At most one of the following statements is true. If one is, circle it. Otherwise, circle "NONE OF THE ABOVE".

- (a) f is increasing on $(-\infty, 1]$, decreasing on $[1, 2]$ and increasing on $[2, \infty)$.
- (b) f is decreasing on $(-\infty, 1]$, increasing on $[1, 2]$ and decreasing on $[2, \infty)$.
- (c) f is increasing on $(-\infty, -2]$, decreasing on $[-2, -1]$ and increasing on $[-1, \infty)$.
- (d) f is decreasing on $(-\infty, -2]$, increasing on $[-2, -1]$ and decreasing on $[-1, \infty)$.
- (e) NONE OF THE ABOVE

$$\begin{aligned} f'(x) &= -(x^2 - 3x + 2) \\ &= -(x-1)(x-2) \end{aligned}$$



II. True or false (no partial credit):

a. (5 pts) Every global extremum occurs at a critical number.

T

b. (5 pts) Every local extremum occurs at a critical number.

T

c. (5 pts) If $f'(3) = 0$ and $f''(3) > 0$, then f has a local maximum at 3.

U

F

d. (5 pts) If f and g are differentiable, then $\frac{d}{dx}[(f(x))(g(x))] = [f'(x)][g'(x)]$.

F

e. (5 pts) If f is increasing on an interval I , then $f' > 0$ on I .

~

F

THE BOTTOM OF THIS PAGE IS FOR TOTALING SCORES
PLEASE DO NOT WRITE BELOW THE LINE

VERSION D

I. A,B,C

I. D,E,F

II. a,b,c,d,e

III. 1.

III. 2.

III. 3,4.

III. 5.

III. Computations. Show work. Unless otherwise specified, answers must be exactly correct, but can be left in any form easily calculated on a standard calculator.

1. (10 pts) Compute $\frac{d}{dx} \left[\frac{2x^3 - 8}{\arctan x} + xe^{\sin x} \right]$

$$\left[\frac{(\arctan x)(6x^2) - (2x^3 - 8)\left(\frac{1}{1+x^2}\right)}{(\arctan x)^2} \right] +$$

$$\left[e^{\sin x} + x(e^{\sin x})(\cos x) \right]$$

2. (10 pts) Using implicit differentiation (and logarithmic differentiation), find $y' = dy/dx$, assuming that $(2 + y^2)^{xy} = 9$.

$$\cancel{[(2+y^2)^{xy}]} \left[\frac{d}{dx} [xy (\ln(2+y^2))] \right] = 0$$

$$y(\ln(2+y^2)) + xy'(\ln(2+y^2)) + xy \left(\frac{2yy'}{2+y^2} \right) = 0$$

$$y' = \frac{-y(\ln(2+y^2))}{x(\ln(2+y^2)) + \frac{2xy^2}{2+y^2}}$$

5. (10 pts) Among all pairs of positive numbers x and y such that $xy = 100$, find the global maximum value of $x + 4y$, provided it exists. Then find the global minimum value, provided it exists. (NOTE: If the global maximum value does not exist, you need to state that clearly to receive full credit. If it does exist, for full credit, you'll need to compute $x + 4y$; computing x and/or y alone is insufficient. These same comments apply to the global minimum value.)

$$y = \frac{100}{x}$$

$$\text{Let } f(x) = x + 4y = x + \frac{400}{x} = x + 400x^{-1}$$

$$f'(x) = 1 - 400x^{-2} = 1 - \frac{400}{x^2}$$

$$f' \quad \begin{array}{cccc} & u & \text{neg} & 0 & \text{pos} \\ & | & & | & \\ \hline & 0 & & 20 & \end{array}$$

On $x > 0$,

$f(x)$ has no global maximum

and has one global minimum

$$\text{at } x = 20, y = \frac{100}{20} = 5$$

with global minimum value

$$20 + 4 \cdot 5 = 40$$


3. (5 pts) Suppose f is 1-1 and $g = f^{-1}$ is the inverse of f . Suppose $f(3) = 4$ and $f'(3) = 58$. Compute $g(4)$ and $g'(4)$.

$$g(4) = 3$$

$$g'(4) = \frac{1}{58}$$

4. (10 pts) Find the maximal intervals of increase and decrease for $f(x) = x^3 - 6x^2 + 5$.

$$\begin{aligned} f'(x) &= 3x^2 - 12x \\ &= 3x(x - 4) \end{aligned}$$



f is increasing on $(-\infty, 0]$

decreasing on $[0, 4]$

increasing on $[4, \infty)$