MATH 1271 Fall 2011, Midterm #2 Handout date: Thursday 10 November 2011

PRINT YOUR NAME:

PRINT YOUR TA'S NAME:

WHAT SECTION ARE YOU IN?

Closed book, closed notes, no calculators/PDAs; no reference materials of any kind. Turn off all handheld devices, including cell phones.

Show work; a correct answer, by itself, may be insufficient for credit. Arithmetic need not be simplified, unless the problem requests it.

I understand the above, and I understand that cheating has severe consequences, from a failing grade to expulsion.

SIGN YOUR NAME:

I. Multiple choice

A. (5 pts) (no partial credit) Find the logarithmic derivative of $(2 + \sin x)^x$ w.r.t. x.

- (a) $\cos x$
- (b) $\ln(\cos x)$

(c)
$$(\ln(2+\sin x)) + \left(\frac{x\cos x}{2+\sin x}\right)$$

(d)
$$[(2 + \sin x)^x] \left[(\ln(2 + \sin x)) + \left(\frac{x \cos x}{2 + \sin x} \right) \right]$$

(e) NONE OF THE ABOVE

B. (5 pts) (no partial credit) Find the derivative of $(2 + \sin x)^x$ w.r.t. x.

- (a) $\cos x$
- (b) $\ln(\cos x)$

(c)
$$(\ln(2+\sin x)) + \left(\frac{x\cos x}{2+\sin x}\right)$$

(d)
$$[(2 + \sin x)^x] \left[(\ln(2 + \sin x)) + \left(\frac{x \cos x}{2 + \sin x} \right) \right]$$

(e) NONE OF THE ABOVE

C. (5 pts) (no partial credit) Compute $\lim_{x\to 0} \left[\frac{\sin^2 x}{4x^3 + 2x^2} \right]$.

- (a) 2
- (b) 1
- (c) 1/2
- (d) 1/4
- (e) NONE OF THE ABOVE

D. (5 pts) (no partial credit) Find the slope of the tangent line to $y = (x^3 + 4)e^{2x}$ at the point (0,4).

- (a) 2
- (b) 4
- (c) 6
- (d) 8
- (e) NONE OF THE ABOVE

E. (5 pts) (no partial credit) Find the logarithmic derivative of $x^2 + 3x - 8$ w.r.t. x.

(a)
$$\frac{2x+3}{x^2+3x-8}$$

(b)
$$\frac{x^2 + 3x - 8}{2x + 3}$$

(c)
$$(\ln(x^2)) + 3(\ln x) - (\ln 8)$$

- (d) ln(2x+3)
- (e) NONE OF THE ABOVE

F. (5 pts) (no partial credit) Suppose $f'(x) = -x^2 + 3x - 2$. At most one of the following statements is true. If one is, circle it. Otherwise, circle "NONE OF THE ABOVE".

- (a) f is increasing on $(-\infty, 1]$, decreasing on [1, 2] and increasing on $[2, \infty)$.
- (b) f is decreasing on $(-\infty, 1]$, increasing on [1, 2] and decreasing on $[2, \infty)$.
- (c) f is increasing on $(-\infty, -2]$, decreasing on [-2, -1] and increasing on $[-1, \infty)$.
- (d) f is decreasing on $(-\infty, -2]$, increasing on [-2, -1] and decreasing on $[-1, \infty)$.
- (e) NONE OF THE ABOVE

- II. True or false (no partial credit):
- a. (5 pts) Every global extremum occurs at a critical number.
- b. (5 pts) Every local extremum occurs at a critical number.
- c. (5 pts) If f'(3) = 0 and f''(3) > 0, then f has a local maximum at 3.
- d. (5 pts) If f and g are differentiable, then $\frac{d}{dx}[(f(x))(g(x))] = [f'(x)][g'(x)]$.
- e. (5 pts) If f is increasing on an interval I, then f' > 0 on I.

THE BOTTOM OF THIS PAGE IS FOR TOTALING SCORES PLEASE DO NOT WRITE BELOW THE LINE

VERSION D

- I. A,B,C
- I. D,E,F
- II. a,b,c,d,e
- III. 1.
- III. 2.
- III. 3,4.
- III. 5.

III. Computations. Show work. Unless otherwise specified, answers must be exactly correct, but can be left in any form easily calculated on a standard calculator.

1. (10 pts) Compute
$$\frac{d}{dx} \left[\frac{2x^3 - 8}{\arctan x} + xe^{\sin x} \right]$$

2. (10 pts) Using implicit differentiation (and logarithmic differentiation), find y'=dy/dx, assuming that $(2+y^2)^{xy}=9$.

3. (5 pts) Suppose f is 1-1 and $g = f^{-1}$ is the inverse of f. Suppose f(3) = 4 and f'(3) = 58. Compute g(4) and g'(4).

4. (10 pts) Find the maximal intervals of increase and decrease for $f(x) = x^3 - 6x^2 + 5$.

5. (10 pts) Among all pairs of positive numbers x and y such that xy = 100, find the global maximum value of x + 4y, provided it exists. Then find the global minimum value, provided it exists. (NOTE: If the global maximum value does not exist, you need to state that clearly to receive full credit. If it does exist, for full credit, you'll need to compute x + 4y; computing x and/or y alone is insufficient. These same comments apply to the global minimum value.)