

(July 18, 2011)

Vector-Valued Integrals

Paul Garrett garrett@math.umn.edu http://www.math.umn.edu/~garrett/

1. Gelfand-Pettis integrals
2. Coproducts, colimits of topological vectorspaces
3. Ubiquity of quasi-complete spaces
4. Totally bounded sets in topological vectorspaces
5. Quasi-completeness and convex hulls of compacts
6. Existence of integrals
7. Historical notes and references

We describe a useful class of topological vectorspaces^[1] V so that *continuous compactly-supported* V -valued functions have *integrals* with respect to finite Borel measures. Rather than *constructing* integrals as limits following [Bochner 1935], [Birkhoff 1935], *et alia*, we use the [Gelfand 1936]-[Pettis 1938] *characterization* of integrals, which has good functorial properties and gives a forceful categorical reason for uniqueness. The issue is *existence*.

A convenient property of topological vectorspaces guaranteeing existence of Gelfand-Pettis integrals is *quasi-completeness*, discussed below. Hilbert, Banach, Fréchet, and LF spaces fall in this class, as do their weak-star duals, and other spaces of mappings such as the *strong operator topology* on mappings between Hilbert spaces, in addition to the *uniform* operator topology.

A compelling application of this integration theory is to holomorphic vector-valued functions, with well-known application to the resolvents of operators on Hilbert and Banach spaces, as in [Dunford 1938] and [Taylor 1938]. In these sources Liouville's theorem on bounded entire \mathbb{C} -valued functions is invoked to prove that a bounded operator on a Hilbert or Banach spaces has non-empty spectrum. A more sophisticated application is to meromorphically-continued Eisenstein series.

Another sort of application of holomorphic and meromorphic vector-valued functions is to *generalized functions*, as in [Gelfand-Shilov 1964], studying *holomorphically parametrized families* of distributions. Many distributions which are not classical functions appear naturally as residues or analytic continuations of families of classical functions with a complex parameter.

A good theory of integration allows a natural treatment of convolutions of distributions with test functions, and related operations.

Two basic auxiliary results are proven about bounded sets in topological vector spaces, the first attributed in [Horvath 1966] to Dieudonné-Schwartz. This result also appears in [Bourbaki 1987], III.5. The second is preparation to invoke a form of the Banach-Steinhaus theorem in situations where the Baire category argument is not directly applicable.

1. Gelfand-Pettis integrals

Let V be a complex topological vectorspace. Let f be a measurable V -valued function on a measure space X . A **Gelfand-Pettis integral** of f would be a vector $I_f \in V$ so that,

$$\lambda(I_f) = \int_X \lambda \circ f \quad (\text{for all } \lambda \in V^*)$$

[1] All vectorspaces here are *complex*, or possibly *real*, as opposed to p -adic or other possibilities.

Assuming that it exists and is unique, this vector I_f would be denoted by

$$I_f = \int_X f$$

By contrast to *construction* of integrals as limits of finite sums, this definition gives a *property* that no reasonable notion of integral would lack, without asking how the property comes to be. Since this property is an irreducible minimum, this definition of integral is called a **weak integral**.

Uniqueness of the integral is immediate when V^* separates points, as for *locally convex* spaces by Hahn-Banach. Similarly, *linearity* of $f \rightarrow I_f$ follows when V^* separates points. Thus, the issue is *existence*. [2]

The functions we integrate are relatively nice: compactly-supported and continuous, on measure spaces with *finite, positive, Borel* measures. In this situation, all the \mathbb{C} -valued integrals

$$\int_X \lambda \circ f$$

exist for elementary reasons, being integrals of compactly-supported \mathbb{C} -valued continuous functions on a compact set with respect to a finite Borel measure.

The technical condition on the topological vectorspace V is that *the convex hull of a compact set must have compact closure*. A more intuitive property which *implies* this property is **quasi-completeness** (and local convexity), meaning that *bounded Cauchy nets* converge. [3] In all applications, when the compactness of closures of convex hulls of compacts holds, it seems that the space is quasi-complete. Thus, while *a priori* the condition of quasi-completeness is stronger than the compactness condition, no example of a strict comparison seems immediate.

The hypotheses of the following theorem will be verified for Fréchet spaces and their weak star duals, for example. In addition to Hilbert and Banach spaces, this includes *Schwartz functions* and *tempered distributions*.

Further, the hypothesis will include *LF-spaces* and their weak $*$ -duals. By definition, an LF space is a countable ascending union of Fréchet spaces so that each Fréchet subspace is closed in the next, with the finest possible topology on the union. These are *strict inductive limits* or *strict colimits* of Fréchet spaces. This includes spaces of *test functions*. The weak $*$ -duals of the LF spaces are also quasi-complete, so we can integrate *distribution-valued* functions.

In addition to the *uniform* operator topology on continuous linear maps from one Hilbert space or Banach space to another, quasi-completeness holds for the *strong* and *weak* operator topologies. [4]

[1.0.1] **Theorem:** Let X be a locally compact Hausdorff topological space with a *finite, positive, Borel* measure. Let V be a locally convex topological vectorspace in which the *closure of the convex hull of a*

[2] Lest there be any doubt, we do require that the integral of a V -valued function be a vector in the space V itself, rather than in a larger space containing V , such as a double dual V^{**} , for example. Some alternative discussions of integration do allow integrals to exist in larger spaces.

[3] In topological vectorspaces which may not have countable local bases, quasi-completeness is more relevant than ‘plain’ completeness. For example, the weak $*$ -dual of an infinite-dimensional Hilbert space is *never* complete, but is always quasi-complete. This example is non-trivial, but helps illustrate the appropriateness of quasi-completeness.

[4] The strong and weak operator topologies, as opposed to the uniform norm topology, are the topologies on operators found useful in representation theory.

compact set is compact. It suffices that V be *quasi-complete*. Then *continuous compactly-supported* V -valued functions f on X have Gelfand-Pettis integrals. Further,

$$\int_X f \in \text{meas}(X) \cdot \left(\text{closure of convex hull of } f(X) \right) \quad (\text{Proof later.})$$

[1.0.2] **Remark:** The conclusion that the integral of f lies in the closure of a convex hull, is a substitute for the estimate of a \mathbb{C} -valued integral by the integral of its absolute value.

2. Coproducts, colimits of topological vectorspaces

The intuitive idea of a topological vectorspace being an *ascending union* of subspaces is necessary, but must be made precise. Spaces of *test functions* on \mathbb{R}^n are obvious prototypes, being ascending unions of Fréchet spaces without being Fréchet spaces themselves. [5]

[2.1] **A categorical viewpoint** *Nested intersections* are examples of *limits*, and limits are *closed subobjects* of the corresponding *products*, while *ascending unions* are examples of *colimits*, and colimits are *quotients* (by closed subobjects) of the corresponding *coproducts*. [6] Thus, proof of existence of *products* gets us half-way to proof of existence of *limits*, and proof of existence of *coproducts* gets us half-way to existence of *colimits*.

In more detail: locally convex *products* of locally convex topological vector spaces are proved to *exist* by constructing them as products of topological spaces, with the product topology, with scalar multiplication and vector addition *induced*.

Given the existence of locally convex *products*, a *limit* of topological vector spaces X_α with compatibility maps $p_\beta^\alpha : X_\alpha \rightarrow X_\beta$ is proven to exist by constructing it as the (closed) subobject Y of the product X of the X_α consisting of $x \in X$ meeting the compatibility conditions

$$p_\beta(x) = (p_\beta^\alpha \circ p_\alpha)(x) \quad (\text{for all } \alpha < \beta, \text{ where } p_\ell \text{ is the projection } X \rightarrow X_\ell)$$

Since local convexity is preserved by these constructions, these constructions do take place inside the category of *locally convex* topological vector spaces.

[2.2] **Coproducts and colimits** *Locally convex* coproducts X of topological vector spaces X_α are coproducts of the vector spaces X_α with the *diamond topology*, described as follows. [7] For a collection U_α of convex neighborhoods of 0 in the X_α , let

$$U = \text{convex hull in } X \text{ of the union of } j_\alpha(U_\alpha) \quad (\text{with } j_\alpha : X_\alpha \rightarrow X \text{ the } \alpha^{\text{th}} \text{ canonical map})$$

[5] A countable ascending union of complete metric spaces, with each a proper closed subspace of the next, *cannot* be complete metric, because it is *presented* as a countable union of nowhere-dense closed subsets, contradicting the conclusion of the Baire Category Theorem.

[6] Categorical limits are often called *projective* limits, and *colimits* are often called *inductive* or *direct* limits. For our purposes, the terms *limit* and *colimit* suffice.

[7] The product topology of locally convex topological vector spaces is unavoidably locally convex, whether the product is in the category of locally convex topological vector spaces or in the larger category of not-necessarily-locally-convex topological vector spaces. However, coproducts behave differently: the locally convex coproduct of *uncountably many* locally convex spaces is *not* a coproduct in the larger category of not-necessarily-locally-convex spaces. This already occurs with an uncountable coproduct of *lines*.

The diamond topology has local basis at 0 consisting of such U . Thus, it is locally convex by construction. (The closedness of points follows from the corresponding property of the X_α .) Thus, existence of a *locally convex* coproduct (of locally convex spaces) is assured by the construction. The *locally convex* colimit of the X_α with compatibility maps $j_\beta^\alpha : X_\alpha \rightarrow X_\beta$ is the *quotient* of the locally convex coproduct X of the X_α by the *closure* of the subobject Z spanned by vectors

$$j_\alpha(x_\alpha) - (j_\beta \circ j_\beta^\alpha)(x_\alpha) \quad (\text{for all } \alpha < \beta \text{ and } x_\alpha \in X_\alpha)$$

Annihilation of these differences in the quotient forces the desired compatibility relations. Obviously, quotients of locally convex spaces are locally convex.

[2.3] **Strict colimits** Let V be a locally convex topological vectorspace with subspaces

$$V_0 \subset V_1 \subset V_2 \subset V_3 \subset \dots \quad (\text{proper containments, } V_i \text{ closed in } V_{i+1})$$

and

$$V = \bigcup_i V_i = \text{colim}_i V_i$$

Such V is a **strict (locally convex) colimit** or **strict (locally convex) inductive limit** of the V_i , *strict* because V_i is closed in V_{i+1} . The fact that the collection of subspaces is *countable* and *well-ordered* is also a special feature. We noted above that locally convex colimits *exist*.

Because of the strictness, we can show that each V_i *injects* to V , as follows. By Hahn-Banach and induction, the identity map $T_i : V_i \rightarrow V_i$ extends compatibly to $T_\beta : V_\ell \rightarrow V_i$ for all ℓ , then yielding a unique compatible $T : V \rightarrow V_i$. The compatibility

$$T \circ j_i = T_i = \text{identity on } V_i \quad (\text{with } j_i : V_i \rightarrow V)$$

implies that j_i is injective. Similarly, each V_i is *closed* in V , as follows. For $v \in V$ but $v \notin V_i$, there is $\ell > i$ such that $v \in V_\ell$, and the quotient map

$$q_\ell : V_\ell \longrightarrow V_\ell/V_i$$

is a map of topological vector spaces (since V_i is closed in V_ℓ) and does not map v to 0. Using Hahn-Banach and well-ordering, q_ℓ extends to compatible maps $q_k : V_k \rightarrow V_k/V_i$ for all $k > i$, and, thus, to a unique compatible map $q : V \rightarrow V_\ell/V_i$. The compatibility assures that $q(v) \neq 0$. That is, v could not be in the closure of V_i , or else it could not be mapped to something non-zero by a continuous topological vector space map that maps V_i to $\{0\}$.

[2.3.1] **Proposition:** Assume that V is a *locally convex strict colimit* of a countable well-ordered collection of closed subspaces V_i . A subset B of V is *bounded* if and only if it lies inside some subspace V_i and is a bounded subset of V_i .

Proof: Suppose B does *not* lie in any V_i . Then there is a sequence i_1, i_2, \dots of positive integers and x_{i_ℓ} in $V_{i_\ell} \cap B$ with x_{i_ℓ} *not* lying in $V_{i_\ell-1}$. Using the simple nature of the indexing set and the simple inter-relationships of the subspaces V_i ,

$$V = \bigcup_j V_{i_\ell}$$

In particular, without loss of generality, we may suppose that $i_\ell = \ell$.

By the Hahn-Banach theorem and induction, using the closedness of V_{i-1} in V_i , there are continuous linear functionals λ_i on V_i 's such that $\lambda_i(x_i) = i$ and the restriction of λ_i to V_{i-1} is λ_{i-1} , for example. Since V is the colimit of the V_i , this collection of functionals exactly describes a unique compatible continuous linear functional λ on V .

But $\lambda(B)$ is *bounded* since B is bounded and λ is continuous, which precludes the possibility that λ takes on all positive integer values at the points x_i of B . Thus, it could *not* have been that B failed to lie inside some single V_i . The strictness of the colimit implies that B is bounded as a subset of V_i , proving one direction of the equivalence. The other direction of the equivalence is less interesting. ///

[2.4] **Equicontinuity on strict colimits** Recall that a set E of continuous linear maps from one topological vectorspace X to another topological vectorspace Y is **equicontinuous** when, for every neighborhood U of 0 in Y , there is a neighborhood N of 0 in X so that $T(N) \subset U$ for every $T \in E$.

[2.4.1] **Proposition:** Let V be a strict colimit of a well-ordered countable collection of locally convex closed subspaces V_i . Let Y be a locally convex topological vectorspace. Let E be a set of continuous linear maps from V to Y . Then E is *equicontinuous* if and only if for each index i the collection of continuous linear maps $\{T|_{V_i} : T \in E\}$ is equicontinuous.

Proof: Given a neighborhood U of 0 in Y , shrink U if necessary so that U is convex and balanced. For each index i , let N_i be a convex, balanced neighborhood of 0 in V_i so that $T(N_i) \subset U$ for all $T \in E$. Let N be the convex hull of the union of the N_i . By the convexity of N , still $T(N) \subset U$ for all $T \in E$. By the construction of the diamond topology, N is an open neighborhood of 0 in the coproduct, hence in the colimit, giving the equicontinuity of E . The other direction of the implication is easy. ///

3. Ubiquity of quasi-complete spaces

This section shows that quasi-completeness is preserved by various important constructions on topological vector spaces, so that most topological vector spaces of interest are quasi-complete. Thus, the theorem on Gelfand-Pettis integrals applies to all these.

Again, a topological vectorspace is **quasi-complete** if every *bounded* Cauchy net is *convergent*. Note that without the assumption that there is a countable local basis (at 0) it is necessary to consider *nets* rather than simply *sequences*. Since many spaces of interest occurring as weak star duals certainly fail to be locally countable, this distinction is not frivolous.

Certainly Fréchet spaces (hence Hilbert and Banach spaces) are quasi-complete, since in the case of a metric space quasi-completeness and ordinary completeness are identical. ^[8]

It is clear that *closed subspaces* of quasi-complete spaces are quasi-complete. Products (with product topology) and finite sums of quasi-complete spaces are quasi-complete.

[3.0.1] **Proposition:** A strict colimit of a countable collection of closed quasi-complete spaces is quasi-complete.

Proof: We saw that bounded subsets of such colimits are exactly the bounded subsets of the limitands. Thus, bounded Cauchy nets in the colimit must be bounded Cauchy nets in one of the closed subspaces. Each of these is assumed quasi-complete, so the colimit is quasi-complete. ///

As a consequence of the proposition, spaces of test functions are quasi-complete, since they are such colimits of the Fréchet spaces of spaces of test functions with prescribed compact support.

Let $\text{Hom}^o(X, Y)$ be the space of continuous linear functions from a topological vectorspace X to another topological vectorspace Y . Give $\text{Hom}^o(X, Y)$ the topology induced by the seminorms $p_{x,U}$ where $x \in X$ and U is a convex, balanced neighborhood of 0 in Y , defined by

$$p_{x,U}(T) = \inf \{t > 0 : Tx \in tU\} \quad (\text{for } T \in \text{Hom}^o(X, Y))$$

[8] That sequential completeness implies completeness for metric spaces is an interesting foundational exercise.

[3.0.2] **Remark:** In the case that X and Y are Hilbert spaces, this construction gives the *strong operator topology* on $\text{Hom}^o(X, Y)$. Replacing the topology on the Hilbert space Y by its *weak* topology, the construction gives $\text{Hom}^o(X, Y)$ the *weak operator topology*. That the collection of continuous linear operators is the same in both cases is a consequence of the Banach-Steinhaus theorem, which also plays a role in the following result.

[3.0.3] **Theorem:** When X is a Fréchet space or LF space, and when Y is quasi-complete, the space $\text{Hom}^o(X, Y)$, with the topology induced by the seminorms $p_{x,U}$ where $x \in X$ and U is a convex, balanced neighborhood of 0 in Y , is *quasi-complete*.

[3.0.4] **Remark:** In fact, the starkest hypothesis on $\text{Hom}^o(X, Y)$ is simply that it support the conclusion of the Banach-Steinhaus theorem. That is, a subset E of $\text{Hom}^o(X, Y)$ so that the set of all images

$$Ex = \{Tx : T \in E\}$$

is bounded (in Y) for all $x \in X$ is necessarily *equicontinuous*. When X is a Fréchet space, this is true (by the usual Banach-Steinhaus theorem) for *any* Y . Further, by the result above on bounded subsets of special sorts of colimits, we see that the same conclusion holds for X such a colimit.

Proof: Let $E = \{T_i : i \in I\}$ be a bounded Cauchy net in $\text{Hom}^o(X, Y)$, where I is a directed set. Of course, attempt to define the limit of the net by

$$Tx = \lim_i T_i x$$

For $x \in X$ the evaluation map $S \rightarrow Sx$ from $\text{Hom}^o(X, Y)$ to Y is continuous. In fact, the topology on $\text{Hom}^o(X, Y)$ is the coarsest with this property. Therefore, by the quasi-completeness of Y , for each fixed $x \in X$ the net $T_i x$ in Y is bounded and Cauchy, so converges to an element of Y suggestively denoted Tx .

To prove *linearity* of T , fix x_1, x_2 in X , $a, b \in \mathbb{C}$ and fix a neighborhood U_o of 0 in Y . Since T is in the closure of E , for any open neighborhood N of 0 in $\text{Hom}^o(X, Y)$, there exists

$$T_i \in E \cap (T + N)$$

In particular, for any neighborhood U of 0 in Y , take

$$N = \{S \in \text{Hom}^o(X, Y) : S(ax_1 + bx_2) \in U, S(x_1) \in U, S(x_2) \in U\}$$

Then

$$\begin{aligned} & T(ax_1 + bx_2) - aT(x_1) - bT(x_2) \\ &= (T(ax_1 + bx_2) - aT(x_1) - bT(x_2)) - (T_i(ax_1 + bx_2) - aT_i(x_1) - bT_i(x_2)) \end{aligned}$$

since T_i is linear. The latter expression is

$$\begin{aligned} & T(ax_1 + bx_2) - (ax_1 + bx_2) + a(T(x_1) - T_i(x_1)) + b(T(x_2) - T_i(x_2)) \\ & \in U + aU + bU \end{aligned}$$

By choosing U small enough so that

$$U + aU + bU \subset U_o$$

we find that

$$T(ax_1 + bx_2) - aT(x_1) - bT(x_2) \in U_o$$

Since this is true for every neighborhood U_o of 0 in Y ,

$$T(ax_1 + bx_2) - aT(x_1) - bT(x_2) = 0$$

which proves linearity.

To prove *continuity* of the limit operator T , we must first be sure that E is *equicontinuous*. For each $x \in X_j$, the set $\{T_i x : i \in I\}$ is bounded in Y , so by Banach-Steinhaus $\{T_i : i \in I\}$ is an equicontinuous set of linear maps from X_i to Y . (Each X_i is Fréchet.) From the result of the previous section on equicontinuous subsets of LF spaces, E itself is equicontinuous.

Fix a neighborhood U of 0 in Y . Invoking the equicontinuity of E , let N be a small enough neighborhood of 0 in X so that $T(N) \subset U$ for all $T \in E$. Let $x \in N$. Choose an index i sufficiently large so that $Tx - T_i x \in U$, vis the definition of the topology on $\text{Hom}^o(X, Y)$. Then

$$Tx \in U + T_i x \in U + U$$

The usual rewriting, replacing U by U' such that $U' + U' \subset U$, shows that T is continuous. ///

4. *Totally bounded sets in topological vectorspaces*

The point of this section is the last corollary, that *convex hulls of compact sets in Fréchet spaces have compact closures*. This is the key point for existence of Gelfand-Pettis integrals.

In preparation, we review the relatively elementary notion of *totally bounded subset* of a metric space, as well as the subtler notion of *totally bounded subset* of a topological vectorspace.

A subset E of a *complete metric space* X is **totally bounded** if, for every $\varepsilon > 0$ there is a covering of E by *finitely-many* open balls of radius ε . The property of *total boundedness* in a metric space is generally stronger than mere *boundedness*. It is immediate that any subset of a totally bounded set is totally bounded.

[4.0.1] Proposition: A subset of a complete metric space has compact closure if and only if it is *totally bounded*.

Proof: Certainly if a set has compact closure then it admits a finite covering by open balls of arbitrarily small (positive) radius.

On the other hand, suppose that a set E is totally bounded in a complete metric space X . To show that E has compact closure it suffices to show that any sequence $\{x_i\}$ in E has a Cauchy subsequence.

We choose such a subsequence as follows. Cover E by finitely-many open balls of radius 1. In at least one of these balls there are infinitely-many elements from the sequence. Pick such a ball B_1 , and let i_1 be the smallest index so that x_{i_1} lies in this ball.

The set $E \cap B_1$ is still totally bounded (and contains infinitely-many elements from the sequence). Cover it by finitely-many open balls of radius $1/2$, and choose a ball B_2 with infinitely-many elements of the sequence lying in $E \cap B_1 \cap B_2$. Choose the index i_2 to be the smallest one so that both $i_2 > i_1$ and so that x_{i_2} lies inside $E \cap B_1 \cap B_2$.

Proceeding inductively, suppose that indices $i_1 < \dots < i_n$ have been chosen, and balls B_i of radius $1/i$, so that

$$x_i \in E \cap B_1 \cap B_2 \cap \dots \cap B_i$$

Then cover $E \cap B_1 \cap \dots \cap B_n$ by finitely-many balls of radius $1/(n+1)$ and choose one, call it B_{n+1} , containing infinitely-many elements of the sequence. Let i_{n+1} be the first index so that $i_{n+1} > i_n$ and so that

$$x_{i_{n+1}} \in E \cap B_1 \cap \dots \cap B_{n+1}$$

Then for $m < n$ we have

$$d(x_{i_m}, x_{i_n}) \leq \frac{1}{m}$$

so this subsequence is Cauchy. ///

In a *topological vectorspace* V , a subset E is **totally bounded** if, for every neighborhood U of 0 there is a finite subset F of V so that

$$E \subset F + U$$

Here the notation $F + U$ means, as usual,

$$F + U = \bigcup_{v \in F} v + U = \{v + u : v \in F, u \in U\}$$

[4.0.2] **Remark:** In a topological vectorspace whose topology is given by a *translation-invariant* metric, a subset is *totally bounded* in this topological vectorspace sense if and only if it is totally bounded in the metric space sense, from the definitions.

[4.0.3] **Lemma:** In a topological vectorspace the convex hull of a *finite* set is *compact*.

Proof: Let the finite set be $F = \{x_1, \dots, x_n\}$. Let σ be the compact set

$$\sigma = \{(c_1, \dots, c_n) \in \mathbb{R}^n : \sum_i c_i = 1, 0 \leq c_i \leq 1, \text{ for all } i\} \subset \mathbb{R}^n$$

Then the convex hull of F is the continuous image of σ under the map

$$(c_1, \dots, c_n) \rightarrow \sum_i c_i x_i$$

so is compact. ///

[4.0.4] **Proposition:** A totally bounded subset E of a *locally convex* topological vectorspace V has totally bounded *convex hull*.

Proof: Let U be a neighborhood of 0 in V . Let U_1 be a *convex* neighborhood of 0 so that $U_1 + U_1 \subset U$. Then for some finite subset F we have $E \subset F + U_1$, by the total boundedness. Let K be the convex hull of F , which by the previous result is *compact*. Then $E \subset K + U_1$, and the latter set is *convex*, as observed earlier. Therefore, the convex hull H of E lies inside $K + U_1$. Since K is compact, it is totally bounded, so can be covered by a finite union $\Phi + U_1$ of translates of U_1 . Thus, since $U_1 + U_1 \subset U$,

$$H \subset (\Phi + U_1) + U_1 \subset \Phi + U$$

Thus, H lies inside this finite union of translates of U . This holds for any open U containing 0, so H is totally bounded. ///

[4.0.5] **Corollary:** In a Fréchet space, the closure of the convex hull of a compact set is compact.

Proof: A compact set in a Fréchet space (or in any complete metric space) is totally bounded, as recalled above. By the previous result, the convex hull of a totally bounded set in a Fréchet space (or in any locally convex space) is totally bounded. Thus, this convex hull has compact closure, since totally bounded sets in complete metric spaces have compact closure. ///

5. Quasi-completeness and convex hulls of compacts

Again, a topological vectorspace X is **quasi-complete** if every *bounded Cauchy net* converges.

The following proof borrows an idea from the proof of the Banach-Alaoglu theorem. It reduces the general case to the case of Fréchet spaces, treated in the previous section.

[5.0.1] Proposition: In a quasi-complete locally convex topological vectorspace X , the closure C of the convex hull H of a compact set K is *compact*.

Proof: Since X is locally convex, by the Hahn-Banach theorem its topology is given by a collection of seminorms v . For each seminorm v , let X_v be the completion of the quotient

$$X/\{x \in X : v(x) = 0\}$$

with respect to the *metric* that v induces on the latter quotient. Thus, X_v is a Fréchet space. (Indeed, the latter quotient is the largest quotient of X on which v induces a *metric* rather than merely a *pseudometric*. And, in fact, X_v is Banach, but we don't use this.) Consider

$$Z = \prod_v X_v \quad (\text{with product topology})$$

with the natural injection $j : X \rightarrow Z$, and with projection p_v to the v^{th} factor.

By construction, and by definition of the topology given by the seminorms, j is a homeomorphism to its image. That is, X is homeomorphic to the subset jX of Z , given the subspace topology.

The image $p_v jK$ is compact, being a continuous image of a compact subset of X . Since X_v is Fréchet, the convex hull H_v of $p_v jK$ has compact closure C_v . The convex hull jH of jK is contained in the product $\prod_v H_v$ of the convex hulls H_v of the projections $p_v jK$. By Tychonoff's theorem, the product $\prod_v C_v$ is *compact*.

Since jC is contained in the compact set $\prod_v C_v$, to prove that the closure jC of jH in jX is compact, it suffices to prove that jC is closed in Z . Since jC is a subset of the compact set $\prod_v C_v$, it is totally bounded and so is certainly bounded (in Z , hence in $X \approx jX$). By the quasi-completeness, any Cauchy net in jC converges to a point in jC . Since any point in the closure of jC in Z has a Cauchy net in jC converging to it, jC is closed in Z . This finishes the proof that quasi-completeness implies the compactness of closures of compact hulls of compacta. ///

6. Existence of integrals

Now we will prove existence of integrals: assume that in the topological vectorspace V convex hulls of compacta have compact closures, and prove the *existence* of Gelfand-Pettis integrals $\int_X f$ of continuous compactly-supported V -valued functions f for $\text{meas}(X) < +\infty$. Further, we prove that the desired integral lies in the (by hypothesis) *compact set*

$$\left(\text{closure of convex hull of } f(X) \right) \cdot \text{meas}(X)$$

Proof: To simplify, divide by a constant to make X have total measure 1. We may assume that X is *compact* since the support of f is compact. Let H be the closure of the convex hull of $f(X)$ in V , compact by hypothesis. We will show that there is an integral of f inside H .

For a *finite* subset L of V^* , let

$$V_L = \{v \in V : \lambda v = \int_X \lambda \circ f, \forall \lambda \in L\}$$

And let

$$I_L = H \cap V_L$$

Since H is compact and V_L is closed, I_L is *compact*. Certainly

$$I_L \cap I_{L'} = I_{L \cup L'}$$

for two finite subsets L, L' of V^* . Thus, if we prove that all the I_L are *non-empty*, then it will follow that the intersection of *all* these compact sets I_L is non-empty. (This is the so-called *finite intersection property*.) That is, we will have *existence* of the integral.

To prove that each I_L is non-empty for *finite* subsets L of V^* , choose an ordering $\lambda_1, \dots, \lambda_n$ of the elements of L . Make a continuous linear mapping $\Lambda = \Lambda_L$ from V to \mathbb{R}^n by

$$\Lambda(v) = (\lambda_1 v, \dots, \lambda_n v)$$

Since this map is continuous, the image $\Lambda(f(X))$ is compact in \mathbb{R}^n .

For a finite set L of functionals, the integral

$$y = y_L = \int_X \Lambda f(x) d\mu(x)$$

is readily defined by component-wise integration. Suppose that this point y is in the convex hull of $\Lambda(f(X))$. Since Λ_L is linear, $y = \Lambda_L v$ for some v in the convex hull of $f(X)$. Then

$$\Lambda_L v = y = (\dots, \int \lambda_i f(x) d\mu(X), \dots)$$

Thus, the point v lies in I_L as desired. Granting that y lies in the convex hull of $\Lambda_L(f(x))$, we are done.

To prove that $y = y_L$ as above lies in the convex hull of $\Lambda_L(f(X))$, suppose *not*. From the lemma below, in a *finite-dimensional* space the convex hull of a compact set is still compact, *without* having to take closure. Thus, invoking also the finite-dimensional case of the Hahn-Banach theorem, there would be a linear functional η on \mathbb{R}^n so that $\eta y > \eta z$ for all z in this convex hull. That is, letting $y = (y_1, \dots, y_n)$, there would be real c_1, \dots, c_n so that for all (z_1, \dots, z_n) in the convex hull

$$\sum_i c_i z_i < \sum_i c_i y_i$$

In particular, for all $x \in X$

$$\sum_i c_i \lambda_i(f(x)) < \sum_i c_i y_i$$

Integration of both sides of this over X *preserves ordering*, giving the absurd

$$\sum_i c_i y_i < \sum_i c_i y_i$$

Thus, y *does* lie in this convex hull. ///

[6.0.1] Lemma: The convex hull of a compact set K in \mathbb{R}^n is compact. In particular, we have compactness without taking closure.

Proof: We first claim that, for a set E in \mathbb{R}^n and for any x a point in the convex hull of E , there are $n + 1$ points x_0, x_1, \dots, x_n in E of which x is a convex combination.

By induction, to prove the claim it suffices to consider a convex combination $v = c_1 v_1 + \dots + c_N v_N$ of vectors v_i with $N > n + 1$ and show that v is actually a convex combination of $N - 1$ of the v_i . Further, we can suppose without loss of generality that all the coefficients c_i are non-zero.

Define a linear map

$$L : \mathbb{R}^N \longrightarrow \mathbb{R}^n \times \mathbb{R} \quad \text{by} \quad L(x_1, \dots, x_N) \longrightarrow \left(\sum_i x_i v_i, \sum_i x_i \right)$$

By dimension-counting, since $N > n + 1$ the kernel of L must be non-trivial. Let (x_1, \dots, x_N) be a non-zero vector in the kernel.

Since $c_i > 0$ for every index, and since there are only finitely-many indices altogether, there is a constant c so that $|cx_i| \leq c_i$ for every index i , and so that $cx_{i_o} = c_{i_o}$ for at least one index i_o . Then

$$v = v - 0 = \sum_i c_i v_i - c \cdot \sum_i x_i v_i = \sum_i (c_i - cx_i) v_i$$

Since $\sum_i x_i = 0$ this is still a convex combination, and since $cx_{i_o} = c_{i_o}$ at least one coefficient has become zero. This is the induction, which proves the claim.

Using the previous claim, a point v in the convex hull of K is actually a convex combination $c_o v_o + \dots + c_n v_n$ of $n + 1$ points v_o, \dots, v_n of K . Let σ be the compact set (c_o, \dots, c_n) with $0 \leq c_i \leq 1$ and $\sum_i c_i = 1$. The convex hull of K is the image of the compact set

$$\sigma \times K^{n+1}$$

under the continuous map

$$L : (c_o, \dots, c_n) \times (v_o, v_1, \dots, v_n) \longrightarrow \sum_i c_i v_i$$

so is compact. This proves the lemma, finishing the proof of the theorem. ///

7. Historical notes and references

Most investigation and use of integration of vector-valued functions is in the context of *Banach-space*-valued functions. Nevertheless, the idea of [Gelfand 1936] extended and developed by [Pettis 1938] immediately suggests a viewpoint not confined to the Banach-space case. A hint appears in [Rudin 1991].

This is in contrast to many of the more detailed studies and comparisons of varying notions of integral specific to the Banach-space case, such as [Bochner 1935]. A variety of developmental episodes and results in the Banach-space-valued case is surveyed in [Hildebrandt 1953]. Proofs and application of many of these results are given in [Hille-Phillips 1957]. (The first edition, authored by Hille alone, is sparser in this regard.) See also [Brooks 1969] to understand the viewpoint of those times.

One of the few exceptions to the apparent limitation to the Banach-space case is [Phillips 1940]. However, it seems that in the United States after the Second World War consideration of anything fancier than Banach spaces was not popular.

The present pursuit of the issue of quasi-completeness (and compactness of the closure of the convex hull of a compact set) was motivated originally by the discussion in [Rudin 1991], although the latter does not make clear that this condition is fulfilled in more than Fréchet spaces, and does not mention quasi-completeness. Imagining that these ideas must be applicable to distributions, one might cast about for means to prove

the compactness condition, eventually hitting upon the hypothesis of quasi-completeness in conjunction with ideas from the proof of the Banach-Alaoglu theorem. Indeed, in [Bourbaki 1987] it is shown (by apparently different methods) that quasi-completeness implies this compactness condition, although there the application to vector-valued integrals is not mentioned. [Schaeffer-Wolff 1999] is a very readable account of further important ideas in topological vector spaces.

The fact that a bounded subset of a countable strict inductive limit of closed subspaces must actually be a bounded subset of one of the subspaces, easy to prove once conceived, is attributed to Dieudonne and Schwartz in [Horvath 1966]. See also [Bourbaki 1987], III.5 for this result. Pathological behavior of uncountable colimits was evidently first exposed in [Douady 1963].

Evidently *quotients* of quasi-complete spaces (by closed subspaces, of course) *may fail to be quasi-complete*: see [Bourbaki 1987], IV.63 exercise 10 for a construction.

[Birkhoff 1935] G. Birkhoff, *Integration of functions with values in a Banach space*, Trans. AMS **38** (1935), 357-378.

[Bochner 1935] S. Bochner, *Integration von Funktionen deren Werte die Elemente eines Vektorraumes sind*, Fund. Math., vol. 20, 1935, pp. 262-276.

[Bourbaki 1987] N. Bourbaki, *Topological Vector Spaces*, ch. 1-5, Springer-Verlag, 1987.

[Brooks 1969] J.K. Brooks, *Representations of weak and strong integrals in Banach spaces*, Proc. Nat. Acad. Sci. U.S., 1969, 266-270.

[Douady 1963] A. Douady, *Parties compactes d'un espace de fonctions continues a support compact*, C. R. Acad. Sci. Paris 257, 1963, pp. 2788-2791.

[Dunford 1936] N. Dunford, *Integration of abstract functions*, abstract, Bull. AMS **44** (1936), 178.

[Dunford 1938] N. Dunford, *Uniformity in linear spaces*, Trans. AMS 44, 1938, pp. 305-356.

[Gelfand 1936] I. M. Gelfand, *Sur un lemme de la theorie des espaces lineaires*, Comm. Inst. Sci. Math. de Kharkoff, no. 4, vol. 13, 1936, pp. 35-40.

[Gelfand-Shilov 1964] I. M. Gelfand, G. E. Shilov *Generalized Functions*, vol. 1, translated by E. Saletan, Academic Press, 1964.

[Hildebrandt 1953] T. H. Hildebrandt, *Integration in abstract spaces*, Bull. AMS, vol. 59, 1953, pp. 111-139.

[Hille-Phillips 1957] E. Hille with R. Phillips, *Functional Analysis and Semigroups*, AMS Colloquium Publication, second edition, 1957.

[Horvath 1966] J. Horvath, *Topological Vector Spaces and Distributions*, Addison-Wesley, 1966.

[Pettis 1938] B. J. Pettis, *On integration in vector spaces*, Trans. AMS, vol. 44, 1938, pp. 277-304.

[Phillips 1940] R. S. Phillips, *Integration in a convex linear topological space*, Trans. AMS, vol. 47, 1940, pp. 114-145.

[Rudin 1991] W. Rudin, *Functional Analysis*, second edition, McGraw-Hill, 1991.

[Schaeffer-Wolff 1999] H.H. Schaeffer, with M.P. Wolff, *Topological Vector Spaces*, second edition, Springer, 1999.

[Taylor 1938] A. E. Taylor, *The resolvent of a closed transformation*, Bull. AMS, vol. 44, 1938, pp. 70-74.

[Taylor 1971] A. E. Taylor, *Notes on the history of the uses of analyticity in operator theory*, Amer. Math. Monthly, vol. 78, 1971, pp. 331-342.