

(April 4, 2014)

Examples of operators and spectra

Paul Garrett garrett@math.umn.edu <http://www.math.umn.edu/~garrett/>

[This document is http://www.math.umn.edu/~garrett/m/fun/notes_2012-13/06b_examples_spectra.pdf]

1. Generalities on spectra
2. Positive examples
3. Simplest Rellich compactness lemma
4. Cautionary examples: non-normal operators

The usefulness of the notion of spectrum of an operator on a Hilbert space is the analogy to eigenvalues of operators on finite-dimensional spaces. Naturally, things become more complicated in infinite-dimensional vector spaces.

1. Generalities on spectra

It is convenient to know that spectra of continuous operators are *non-empty, compact* subsets of \mathbb{C} .

Knowing this, *every* non-empty compact subset of \mathbb{C} is easily made to appear as the spectrum of a continuous operator, even *normal* ones, as below.

[1.0.1] Proposition: The spectrum $\sigma(T)$ of a continuous linear operator $T : V \rightarrow V$ on a Hilbert space V is *bounded* by the operator norm $|T|_{\text{op}}$.

Proof: For $|\lambda| > |T|_{\text{op}}$, an obvious heuristic suggests an expression for the *resolvent* $R_\lambda = (T - \lambda)^{-1}$:

$$(T - \lambda)^{-1} = -\lambda^{-1} \cdot \left(1 - \frac{T}{\lambda}\right)^{-1} = -\lambda^{-1} \cdot \left(1 + \frac{T}{\lambda} + \left(\frac{T}{\lambda}\right)^2 + \dots\right)$$

The infinite series converges in operator norm for $|T/\lambda|_{\text{op}} < 1$, that is, for $|\lambda| > |T|_{\text{op}}$. Then

$$(T - \lambda) \cdot (-\lambda^{-1}) \cdot \left(1 + \frac{T}{\lambda} + \left(\frac{T}{\lambda}\right)^2 + \dots\right) = 1$$

giving a continuous inverse $(T - \lambda)^{-1}$, so $\lambda \notin \sigma(T)$. ///

[1.0.2] Remark: The same argument applied to T^n shows that $\sigma(T^n)$ is inside the closed ball of radius $|T^n|_{\text{op}}$. By the elementary identity

$$T^n - \lambda^n = (T - \lambda) \cdot (T^{n-1} + T^{n-2}\lambda + \dots + T\lambda^{n-2} + \lambda^{n-2})$$

$(T - \lambda)^{-1}$ exists for $|\lambda^n| > |T^n|_{\text{op}}$, that is, for $|\lambda| > |T^n|_{\text{op}}^{1/n}$. That is, $\sigma(T)$ is inside the closed ball of radius $\inf_{n \geq 1} |T^n|_{\text{op}}^{1/n}$. The latter expression is the *spectral radius* of T . This notion is relevant to *non-normal* operators, such as the *Volterra operator*, whose spectral radius is 0, while its operator norm is much larger.

[1.0.3] Proposition: The spectrum $\sigma(T)$ of a continuous linear operator $T : V \rightarrow V$ on a Hilbert space V is *compact*.

Proof: That $\lambda \notin \sigma(T)$ is that there is a continuous linear operator $(T - \lambda)^{-1}$. We claim that for μ sufficiently close to λ , $(T - \mu)^{-1}$ exists. Indeed, a heuristic suggests an expression for $(T - \mu)^{-1}$ in terms of $(T - \lambda)^{-1}$. The algebra is helpfully simplified by replacing T by $T + \lambda$, so that $\lambda = 0$. With μ near 0 and granting existence of T^{-1} , the heuristic is

$$(T - \mu)^{-1} = (1 - \mu T^{-1})^{-1} \cdot T^{-1} = \left(1 + \mu T^{-1} + (\mu T^{-1})^2 + \dots\right) \cdot T^{-1}$$

The geometric series converges in operator norm for $|\mu T^{-1}|_{\text{op}} < 1$, that is, for $|\mu| < |T^{-1}|_{\text{op}}^{-1}$. Having found the obvious candidate for an inverse,

$$(1 - \mu T^{-1}) \cdot \left(1 + \mu T^{-1} + (\mu T^{-1})^2 + \dots\right) = 1$$

and

$$(T - \mu) \cdot \left(1 + \mu T^{-1} + (\mu T^{-1})^2 + \dots\right) \cdot T^{-1} = 1$$

so there is a continuous linear operator $(T - \mu)^{-1}$, and $\mu \notin \sigma(T)$. Having already proven that $\sigma(T)$ is bounded, it is compact. ///

[1.0.4] Proposition: The spectrum $\sigma(T)$ of a continuous linear operator on a Hilbert space $V \neq \{0\}$ is non-empty.

Proof: The argument reduces the issue to Liouville's theorem from complex analysis, that a bounded entire (holomorphic) function is constant. Further, an entire function that goes to 0 at ∞ is identically 0.

Suppose the resolvent $R_\lambda = (T - \lambda)^{-1}$ is a continuous linear operator for all $\lambda \in \mathbb{C}$. The operator norm is readily estimated for large λ :

$$\begin{aligned} |R_\lambda|_{\text{op}} &= |\lambda|^{-1} \cdot \left|1 + \frac{T}{\lambda} + \left(\frac{T}{\lambda}\right)^2 + \dots\right|_{\text{op}} \\ &\leq |\lambda|^{-1} \cdot \left(1 + \left|\frac{T}{\lambda}\right|_{\text{op}} + \left|\frac{T}{\lambda}\right|_{\text{op}}^2 + \dots\right) = \frac{1}{|\lambda|} \cdot \frac{1}{1 - \frac{|T|_{\text{op}}}{|\lambda|}} \end{aligned}$$

This goes to 0 as $|\lambda| \rightarrow \infty$. *Hilbert's identity* asserts the complex differentiability as operator-valued function:

$$\frac{R_\lambda - R_\mu}{\lambda - \mu} = R_\lambda \cdot \frac{(T - \mu) - (T - \lambda)}{\lambda - \mu} \cdot R_\mu = R_\lambda \cdot R_\mu \rightarrow R_\lambda^2 \quad (\text{as } \mu \rightarrow \lambda)$$

since $\mu \rightarrow R_\mu$ is continuous for large μ , by the same identity:

$$|R_\lambda - R_\mu|_{\text{op}} \leq |\lambda - \mu| \cdot |R_\mu \cdot R_\lambda|_{\text{op}}$$

Thus, the scalar-valued functions $\lambda \rightarrow \langle R_\lambda v, w \rangle$ for $v, w \in V$ are complex-differentiable, and satisfy

$$|\langle R_\lambda v, w \rangle| \leq |R_\lambda v| \cdot |w| \leq |R_\lambda|_{\text{op}} \cdot |v| \cdot |w| \leq \frac{1}{|\lambda|} \cdot \frac{1}{1 - \frac{|T|_{\text{op}}}{|\lambda|}} \cdot |v| \cdot |w|$$

By Liouville, $\langle R_\lambda v, w \rangle = 0$ for all $v, w \in V$, which is impossible. Thus, the spectrum is not empty. ///

[1.0.5] Proposition: The entire spectrum, both point-spectrum and continuous-spectrum, of a self-adjoint operator is a non-empty, compact subset of \mathbb{R} . The entire spectrum of a unitary operator is a non-empty, compact subset of the unit circle.

Proof: For self-adjoint T , we claim that the imaginary part of $\langle (T - \mu)v, v \rangle$ is at least $\langle v, v \rangle$ times the imaginary part of μ . Indeed, $\langle Tv, v \rangle$ is real, since

$$\langle Tv, v \rangle = \langle v, T^*v \rangle = \langle v, Tv \rangle = \overline{\langle Tv, v \rangle}$$

so

$$\langle (T - \mu)v, v \rangle = \langle Tv, v \rangle - \mu \cdot \langle v, v \rangle$$

and

$$|\operatorname{Im}\langle(T - \mu)v, v\rangle| = |\operatorname{Im}\mu| \cdot \langle v, v\rangle$$

and by Cauchy-Schwarz-Bunyakovsky

$$|(T - \mu)v| \cdot |v| \geq |\langle(T - \mu)v, v\rangle| \geq |\operatorname{Im}\mu| \cdot \langle v, v\rangle = |\operatorname{Im}\mu| \cdot |v|^2$$

Dividing by $|v|$,

$$|(T - \mu)v| \geq |\operatorname{Im}\mu| \cdot |v|$$

This inequality shows more than the injectivity of $T - \mu$. Namely, the inequality gives a bound on the operator norm of the inverse $(T - \mu)^{-1}$ defined on the image of $T - \mu$. The image is *dense* since μ is not an eigenvalue and there is no residual spectrum for normal operators T . Thus, the inverse extends by continuity to a continuous linear map defined on the whole Hilbert space. Thus, $T - \mu$ has a continuous linear inverse, and μ is not in the spectrum of T .

For T unitary, $|Tv| = |v|$ for all v implies $T|_{\text{op}} = 1$. Thus, $\sigma(T)$ is contained in the unit disk, by the general bound on spectra in terms of operator norms. From $(T - \lambda)^* = T^* - \bar{\lambda}$, the spectrum of T^* is obtained by complex-conjugating the spectrum of T . Thus, for unitary T , the spectrum of $T^{-1} = T^*$ is also contained in the unit disk. At the same time, the natural

$$T - \lambda = -T \cdot (T^{-1} - \lambda^{-1}) \cdot \lambda$$

gives

$$(T - \lambda)^{-1} = -\lambda^{-1} \cdot (T^{-1} - \lambda^{-1})^{-1} \cdot T^{-1}$$

so $\lambda^{-1} \in \sigma(T^{-1})$ exactly when $\lambda \in \sigma(T)$. Thus, the spectra of both T and $T^* = T^{-1}$ are inside the unit circle. ///

2. Positive examples

This section gives non-pathological examples. Let ℓ^2 be the usual space of square-summable sequences (a_1, a_2, \dots) , with standard orthonormal basis

$$e_j = \underbrace{(0, \dots, 0, 1, 0, \dots)}_{1 \text{ at } j\text{th position}}$$

[2.1] Multiplication operators with specified eigenvalues Given a countable, bounded list of complex numbers λ_j , the operator $T : \ell^2 \rightarrow \ell^2$ by

$$T : (a_1, a_2, \dots) \longrightarrow (\lambda_1 \cdot a_1, \lambda_2 \cdot a_2, \dots)$$

has λ_j -eigenvector the standard basis element e_j . Clearly

$$T^* : (a_1, a_2, a_3, \dots) \longrightarrow (\bar{\lambda}_1 \cdot a_1, \bar{\lambda}_2 \cdot a_2, \bar{\lambda}_3 \cdot a_3, \dots)$$

so T is *normal*, in the sense that $TT^* = T^*T$. To see that there are no *other* eigenvalues, suppose $Tv = \mu \cdot v$ with μ not among the λ_j . Then

$$\mu \cdot \langle v, e_j \rangle = \langle Tv, e_j \rangle = \langle v, T^*e_j \rangle = \langle v, \bar{\lambda}_j e_j \rangle = \lambda_j \cdot \langle v, e_j \rangle$$

Thus, $(\mu - \lambda_j) \cdot \langle v, e_j \rangle = 0$, and $\langle v, e_j \rangle = 0$ for all j . Since e_j form an orthonormal basis, $v = 0$. ///

[2.2] Every compact subset of \mathbb{C} is the spectrum of an operator Grant for the moment a countable dense subset $\{\lambda_j\}$ of a non-empty compact subset^[1] C of \mathbb{C} , and as above let

$$T : (a_1, a_2, a_3, \dots) \longrightarrow (\lambda_1 \cdot a_1, \lambda_2 \cdot a_2, \lambda_3 \cdot a_3, \dots)$$

We saw that there are no further eigenvalues. Since spectra are *closed*, the closure C of $\{\lambda_j\}$ is *contained* in $\sigma(T)$.

It remains to show that the continuous spectrum is no larger than the closure C of the eigenvalues, *in this example*. That is, for $\mu \notin C$, exhibit a continuous linear $(T - \mu)^{-1}$.

For $\mu \notin C$, there is a uniform lower bound $0 < \delta \leq |\mu - \lambda_j|$. That is, $\sup_j |\mu - \lambda_j|^{-1} \leq \delta^{-1}$. Thus, the naturally suggested map

$$(a_1, a_2, \dots) \longrightarrow \left((\lambda_1 - \mu)^{-1} \cdot a_1, (\lambda_2 - \mu)^{-1} \cdot a_2, \dots \right)$$

is a bounded linear map, and gives $(T - \mu)^{-1}$.

[2.3] Two-sided shift has no eigenvalues Let V be the Hilbert space of *two-sided* sequences $(\dots, a_{-1}, a_0, a_1, \dots)$ with natural inner product

$$\langle (\dots, a_{-1}, a_0, a_1, \dots), (\dots, b_{-1}, b_0, b_1, \dots) \rangle = \dots + a_{-1}b_{-1} + a_0b_0 + a_1b_1 + \dots$$

The right and left *two-sided* shift operators are

$$(R \cdot a)_n = a_{n-1} \qquad (L \cdot a)_n = a_{n+1}$$

These operators are mutual adjoints, mutual inverses, so are unitary. Being unitary, their operator norms are 1, so their spectra are non-empty compact subsets of the unit circle.

They have no eigenvalues: indeed, for $Rv = \lambda \cdot v$, if there is any index n with $v_n \neq 0$, then the relation $Rv = \lambda \cdot v$ gives $v_{n+k+1} = \lambda \cdot v_{n+k}$ for $k = 0, 1, 2, \dots$. Since $|\lambda| = 1$, such a vector is not in ℓ^2 .

Nevertheless, we claim that $\lambda \in \sigma(L)$ for every λ with $|\lambda| = 1$, and similarly for R . Indeed, for λ *not* in the spectrum, there is a continuous linear operator $(L - \lambda)^{-1}$, so $|(L - \lambda)v| \geq \delta \cdot |v|$ for some $\delta > 0$. It is easy to make *approximate* eigenvectors for L for any $|\lambda| = 1$: let

$$v^{(\ell)} = (\dots, 0, \dots, 0, 1, \lambda, \lambda^2, \lambda^3, \dots, \lambda^\ell, 0, 0, \dots)$$

Obviously it doesn't matter where the non-zero entries begin. From

$$(L - \lambda)v^{(\ell)} = (\dots, 0, \dots, 0, 1, 0, \dots, 0, \lambda^{\ell+1}, 0, 0, \dots)$$

$|(L - \lambda)v^{(\ell)}| = \sqrt{1 + 1}$, while $|v^{(\ell)}| = \sqrt{\ell + 1}$. Thus, $|(L - \lambda)v^{(\ell)}|/|v^{(\ell)}| \longrightarrow 0$, and there can be no $(L - \lambda)^{-1}$. Thus, every λ on the unit circle is in $\sigma(R)$.

[2.4] Compact multiplication operators on ℓ^2 For a sequence of complex numbers $\lambda_n \rightarrow 0$, we claim that the multiplication operator

$$T : (a_1, a_2, \dots) \longrightarrow (\lambda_1 \cdot a_1, \lambda_2 \cdot a_2, \dots)$$

[1] To make a countable dense subset of C , for $n = 1, 2, \dots$ cover C by finitely-many disks of radius $1/n$, each meeting C , and in each choose a point of C . The union over $n = 1, 2, \dots$ of these finite sets is countable and dense in C .

is compact. We already showed that it has eigenvalues exactly $\lambda_1, \lambda_2, \dots$, and spectrum the closure of $\{\lambda_j\}$. Thus, the spectrum includes 0, but 0 is an *eigenvalue* only when it appears among the λ_j , which may range from 0 times to infinitely-many times.

To prove that the operator is compact, we prove that the image of the unit ball is pre-compact, by showing that it is *totally bounded*. Given $\varepsilon > 0$, take k such that $|\lambda_i| < \varepsilon$ for $i > k$. The ball in \mathbb{C}^k of radius $\max\{|\lambda_j| : j \leq k\}$ is precompact, so has a finite cover by ε -balls, centered at points v^1, \dots, v^N . For $v = (v_1, v_2, \dots)$ with $|v| \leq 1$,

$$Tv = (\lambda_1 v_1, \lambda_2 v_2, \dots, \lambda_k v_k, 0, 0, \dots) + (0, \dots, 0, \lambda_{k+1} v_{k+1}, \lambda_{k+2} v_{k+2}, \dots)$$

With v^j the closest of the v^1, \dots, v^N to $(\lambda_1 v_1, \lambda_2 v_2, \dots, \lambda_k v_k, 0, 0, \dots)$,

$$|Tv - v^j| < \varepsilon + |(0, \dots, 0, \lambda_{k+1} v_{k+1}, \lambda_{k+2} v_{k+2}, \dots)| < \varepsilon + \varepsilon \cdot |(0, \dots, 0, v_{k+1}, v_{k+2}, \dots)| \leq \varepsilon + \varepsilon \cdot |v| \leq 2\varepsilon$$

Thus, the image of the unit ball under T is covered by finitely-many 2ε -balls. ///

[2.5] Multiplication operators on $L^2[a, b]$ For $\varphi \in C^o[a, b]$, we claim that the multiplication operator

$$M_\varphi : L^2[a, b] \longrightarrow L^2[a, b]$$

by

$$M_\varphi f(x) = \varphi(x) \cdot f(x)$$

is *normal*, and has spectrum the image $\varphi[a, b]$ of φ . The eigenvalues are λ such that $\varphi(x) = \lambda$ on a subset of $[a, b]$ of positive measure. The normality is clear, so, beyond eigenvalues, we need only examine continuous spectrum, not residual.

On one hand, if $\varphi(x) = \lambda$ on a set of positive measure, there is an infinite-dimensional sub-space of $L^2[0, 1]$ of functions supported there, and all these are eigenvectors. On the other hand, if $f \neq 0$ in $L^2[0, 1]$ and $\varphi(x) \cdot f(x) = \lambda \cdot f(x)$, even if f is altered on a set of measure 0, it must be that $\varphi(x) = \lambda$ on a set of positive measure.

To understand the continuous spectrum, for $\varphi(x_o) = \lambda$ make *approximate eigenvectors* by taking L^2 functions f supported on $[x_o - \delta, x_o + \delta]$, where $\delta > 0$ is small enough so that $|\varphi(x) - \varphi(x_o)| < \varepsilon$ for $|x - x_o| < \delta$. Then

$$\|(M_\varphi - \lambda)f\|_{L^2}^2 = \int |\varphi(x) - \lambda|^2 \cdot |f(x)|^2 dx \leq \varepsilon^2 \cdot \|f\|_{L^2}^2$$

Thus, $\inf_{f \neq 0} \|(M_\varphi - \lambda)f\|_{L^2} / \|f\|_{L^2} = 0$, so $M_\varphi - \lambda$ is not invertible. If λ is not an eigenvalue, it is in the continuous spectrum. On the other hand, if $\varphi(x) \neq \lambda$, then there is some $\delta > 0$ such that $|\varphi(x) - \lambda| \geq \delta$ for all $x \in [0, 1]$, by the compactness of $[0, 1]$. Then

$$\|(M_\varphi - \lambda)f\|_{L^2}^2 = \int_0^1 |\varphi(x) - \lambda|^2 \cdot |f(x)|^2 dx \geq \int_0^1 \delta^2 \cdot |f(x)|^2 dx = \delta^2 \cdot \|f\|_{L^2}^2$$

Thus, there is a continuous inverse $(M_\varphi - \lambda)$, and λ is *not* in the spectrum.

3. Simplest Rellich compactness lemma

One characterization of the s^{th} Levi-Sobolev space of functions $H^s(A)$ on a product $A = (S^1)^{\times n}$ of circles $S^1 = \mathbb{R}/2\pi\mathbb{Z}$ is as the closure of the function space of *finite* Fourier series with respect to the Levi-Sobolev norm (squared)

$$\left| \sum_{\xi \in \mathbb{Z}^n} c_\xi e^{i\xi \cdot x} \right|_{H^s}^2 = \sum_{\xi \in \mathbb{Z}^n} |c_\xi|^2 \cdot (1 + |\xi|^2)^s \quad (s \in \mathbb{R}, \text{ on finite Fourier series})$$

The standard orthonormal basis for $H^s(A)$ is

$$\frac{1}{(2\pi)^{n/2}} \cdot \frac{e^{i\xi \cdot x}}{(1 + |\xi|^2)^{s/2}} \quad (\text{with } \xi \in \mathbb{Z}^n)$$

By the Plancherel theorem, the map from $L^2(\mathbb{Z}^n)$ (with counting measure) to $L^2(A)$ by

$$\{c_\xi : \xi \in \mathbb{Z}^n\} \longrightarrow \frac{1}{(2\pi)^{n/2}} \sum_{\xi \in \mathbb{Z}^n} c_\xi \frac{e^{i\xi \cdot x}}{(1 + |\xi|^2)^{s/2}}$$

is an isometric isomorphism.

For $s > t$, there is a continuous inclusion $H^s(A) \rightarrow H^t(A)$. In terms of these orthonormal bases, there is a commutative diagram

$$\begin{array}{ccc} L^2(A) & \xrightarrow{T} & L^2(A) \\ \approx \downarrow & & \downarrow \approx \\ H^s(A) & \xrightarrow{\text{inc}} & H^t(A) \end{array}$$

given by

$$\begin{array}{ccc} \{c_\xi\} & \xrightarrow{T} & \left\{ (1 + |\xi|^2)^{\frac{t-s}{2}} \cdot c_\xi \right\} \\ \approx \downarrow & & \downarrow \approx \\ \frac{1}{(2\pi)^{n/2}} \sum_{\xi} c_\xi \frac{e^{i\xi \cdot x}}{(1 + |\xi|^2)^{s/2}} & \xrightarrow{\text{inc}} & \frac{1}{(2\pi)^{n/2}} \sum_{\xi} (1 + |\xi|^2)^{\frac{t-s}{2}} \cdot c_\xi \frac{e^{i\xi \cdot x}}{(1 + |\xi|^2)^{t/2}} \end{array}$$

Since $s > t$, the number $\lambda_\xi = (1 + |\xi|^2)^{\frac{t-s}{2}}$ are bounded by 1, and have unique limit point 0. In particular, $T : L^2(A) \rightarrow L^2(A)$ is *compact*.

Thus, we have the simplest instance of *Rellich's compactness lemma*: the inclusion $H^s(A) \rightarrow H^t(A)$ is *compact* for $s > t$.

4. Cautionary examples: non-normal operators

[4.1] Shift operators on one-sided ℓ^2 We claim the following: The right-shift

$$R : (a_1, a_2, \dots) \longrightarrow (0, a_1, a_2, \dots)$$

and the left-shift

$$L : (a_1, a_2, a_3, \dots) \longrightarrow (a_2, \dots)$$

are mutual adjoints. These operators are not normal, since $L \circ R = 1_{\ell^2}$ but

$$R \circ L : (a_1, a_2, \dots) \longrightarrow (0, a_2, \dots)$$

The eigenvalues of the left-shift L are all complex numbers in the open unit disk in \mathbb{C} . In particular, there is a *continuum* of eigenvalues and eigenvectors, so they *cannot be mutually orthogonal*. The spectrum $\sigma(L)$ is the closed unit disk.

The right-shift R has *no* eigenvalues, has continuous spectrum the unit circle, and residual spectrum the open unit disk with 0 removed.

Indeed, suppose

$$(0, a_1, a_2, \dots) = R(a_1, a_2, \dots) = \lambda \cdot (a_1, a_2, \dots)$$

With n the lowest index such that $a_n \neq 0$, the n^{th} component in the eigenvector relation gives $0 = a_{n-1} = \lambda \cdot a_n$, so $\lambda = 0$. Then, the $(n+1)^{\text{th}}$ component gives $a_n = \lambda \cdot a_{n+1} = 0$, contradiction. This proves that R has *no* eigenvalues.

Oppositely, for $|\lambda| < 1$,

$$L(1, \lambda, \lambda^2, \dots) = (\lambda, \lambda^2, \dots) = \lambda \cdot (1, \lambda, \lambda^2, \dots)$$

so every such λ is an eigenvector for L . On the other hand, for $|\lambda| = 1$, in an eigenvector relation

$$(a_2, \dots) = L(a_1, a_2, \dots) = \lambda \cdot (a_1, a_2, \dots)$$

let n be the smallest index n with $a_n \neq 0$. Then $a_{n+1} = \lambda \cdot a_n$, $a_{n+2} = \lambda \cdot a_{n+1}$, \dots , so

$$(a_1, a_2, \dots) = (0, \dots, 0, a_n, \lambda a_n, \lambda^2 a_n, \dots)$$

But this is not in ℓ^2 for $|\lambda| = 1$ and $a_n \neq 0$, so λ on the unit circle is *not* an eigenvalue.

For $|\lambda| = 1$, we can make *approximate* λ -eigenvectors for L by

$$v^{[N]} = (1, \lambda, \lambda^2, \dots, \lambda^N, 0, 0, \dots)$$

since

$$(L - \lambda)v^{[N]} = (\lambda, \lambda^2, \dots, \lambda^N, 0, 0, 0, \dots) - \lambda \cdot (1, \lambda, \lambda^2, \dots, \lambda^N, 0, 0, \dots) = (0, 0, \dots, 0, 0, \lambda^{N+1}, 0, 0, \dots)$$

Since

$$\frac{|(L - \lambda)v^{[N]}|}{|v^{[N]}|} = \frac{|\lambda|^{N+1}}{(1 + |\lambda|^2 + \dots + |\lambda|^{2N})^{1/2}} = \frac{1}{\sqrt{N+1}} \rightarrow 0$$

there can be no continuous $(L - \lambda)^{-1}$. Thus, λ on the unit circle is in the spectrum, but not in the point spectrum.

That the unit circle is in the spectrum also follows from the observation above that all λ with $|\lambda| < 1$ are eigenvalues, and the fact that the spectrum is *closed*.

The spectrum of L is bounded by the operator norm $|L|_{\text{op}}$, and $|L|_{\text{op}}$ is visibly 1, so is nothing *else* in the spectrum.

To see that the unit circle is the *continuous* spectrum of L , as opposed to *residual*, we show that $L - \lambda$ has dense image for $|\lambda| = 1$. Indeed, for w such that, for all $v \in \ell^2$,

$$0 = \langle (L - \lambda)v, w \rangle = \langle v, (L^* - \bar{\lambda})w \rangle = \langle v, (R - \bar{\lambda})w \rangle$$

we would have $(R - \bar{\lambda})w = 0$. However, we have seen that R has no eigenvalues. Thus, $L - \lambda$ always has dense image, and the unit circle is continuous spectrum for L .

Reversing that discussion, every λ with $|\lambda| < 1$ is in the residual spectrum of R , because such λ is not an eigenvalue, and $R - \lambda$ does *not* have dense image: for w a $\bar{\lambda}$ -eigenvector for L ,

$$\langle (R - \lambda)v, w \rangle = \langle v, (R^* - \bar{\lambda})w \rangle = \langle v, (L - \bar{\lambda})w \rangle = \langle v, 0 \rangle = 0$$

That is, the image $(R - \lambda)\ell^2$ is in the orthogonal complement to the eigenvector w . The same computation shows that the unit circle is *continuous* spectrum for R , because it is *not* eigenvalues for L .

[4.2] Volterra operator We will show that the Volterra operator $Vf(x) = \int_0^x f(t) dt$ on $L^2[0, 1]$ is *compact*, but not self-adjoint, that its spectrum is $\{0\}$, and that it has no eigenvalues.

A relation $Tf = \lambda \cdot f$ for $f \in L^2$ and $\lambda \neq 0$ implies f is *continuous*:

$$|\lambda| \cdot |f(x+h) - f(x)| = |Tf(x+h) - Tf(x)| \leq \int_x^{x+h} 1 \cdot |f(t)| dt \leq |h|^{\frac{1}{2}} \cdot |f|_{L^2}$$

The fundamental theorem of calculus would imply f is continuously differentiable and $\lambda \cdot f' = (Tf)' = f$. Thus, f would be a constant multiple of $e^{x/\lambda}$, by the mean value theorem. However, by Cauchy-Schwarz-Bunyakowsky, for a λ -eigenfunction

$$|\lambda| \cdot |f(x)| \leq |x|^{\frac{1}{2}} \cdot |f|_{L^2}$$

No non-zero multiple of the exponential satisfies this. Thus, there are no eigenvectors for *non-zero* eigenvalues.

For $f \in L^2[0, 1]$ and $Tf = 0 \in L^2[0, 1]$, Tf is almost everywhere 0. Since $x \rightarrow Tg(x)$ is unavoidably *continuous*, $Tf(x)$ is 0 for all x . Thus, for all x, y in the interval,

$$0 = 0 - 0 = Tf(y) - Tf(x) = \int_x^y f(t) dt$$

That is, $x \rightarrow Tf(x)$ is orthogonal in $L^2[0, 1]$ to all characteristic functions of intervals. Finite linear combinations of these are dense in $C^o[0, 1]$ in the L^2 topology, and $C^o[0, 1]$ is dense in $L^2[0, 1]$. Thus $f = 0$, and there are no eigenvectors for the Volterra operator.

To see that T is *compact*, rewrite it as being given by an *integral kernel* $K(x, y)$:

$$Tf(x) = \int_0^x f(y) dy = \int_0^1 K(x, y) f(y) dy \quad \left(\text{with } K(x, y) = \begin{cases} 0 & (\text{for } 0 \leq y < x) \\ 1 & (\text{for } x < y \leq 1) \end{cases} \right)$$

Thus, T is Hilbert-Schmidt, and compact. The adjoint T^* is given by the integral kernel $K^*(x, y) = \overline{K(y, x)}$, visibly different from $K(x, y)$, so T is *not* self-adjoint.

To see that the spectrum is *at most* $\{0\}$, show that the *spectral radius* is 0:

$$\begin{aligned} T^n f(x) &= \int_0^x \int_0^{x_{n-1}} \dots \int_0^{x_2} \int_0^{x_1} f(t) dt dx_1 \dots dx_{n-1} = \int_0^x f(t) \left(\int_t^x \int_t^{x_{n-1}} \dots \int_t^{x_2} dx_1 \dots dx_{n-1} \right) dt \\ &= \int_0^x f(t) \cdot \frac{(x-t)^{n-1}}{(n-1)!} dt \end{aligned}$$

From this, $|T^n|_{\text{op}} \leq \frac{1}{n!}$, and

$$\begin{aligned} \log \lim_{2n} \left(\frac{1}{(2n)!} \right)^{1/2n} &= -\lim_{2n} \frac{1}{2n} \cdot \log(2n)! = -\lim_{2n} \frac{1}{2n} \sum_{1 \leq k \leq 2n} \log k \\ &= -\lim_{2n} \frac{1}{2n} \sum_{1 \leq k \leq n} (\log k + \log(2n - k + 1)) \leq -\lim_{2n} \frac{1}{2n} \sum_{1 \leq k \leq \frac{n}{k}} (\log k + \log(2n - k + 1)) \\ &\leq -\lim_{2n} \frac{1}{2n} \sum_{1 \leq k \leq \frac{n}{k}} \log 2n = -\lim_{2n} \frac{\log 2n}{2} = -\infty \end{aligned}$$

since $k(2n - k) \geq 2n$ for $1 \leq k \leq n$, noting the sign. That is, $\lim_n |T^n|_{\text{op}}^{1/n} = 0$, so the spectral radius is 0. Since the spectrum is non-empty, it must be exactly $\{0\}$.