

(October 24, 2014)

Adjoint, naturality, exactness, small Yoneda lemma

Paul Garrett garrett@math.umn.edu <http://www.math.umn.edu/~garrett/>

[This document is http://www.math.umn.edu/~garrett/m/repns/notes_2014-15/04b_adjoint_exactness.pdf]

The best way to certify left-exactness or right-exactness of an additive functor^[1] is to see that it appears is a *right adjoint* or *left adjoint*. Many familiar functors occur in pairs whose adjointness is *obvious* once observed. MacLane and others have quipped *Everything's an adjoint*.

The proof that left adjoints are right-exact, and that right adjoints are left-exact, uses a small incarnation of *Yoneda's Lemma*, and illustrates the importance of *naturality* of isomorphisms.

Throughout, we consider only \mathbb{Z} -modules, although the arguments apply more generally.

- $\text{Hom}(X, -)$ is left exact
- Adjoint and naturality
- A small Yoneda lemma
- Half-exactness of adjoints

1. $\text{Hom}(X, -)$ is left exact

Everything later will reduce to the straightforward left-exactness of $\text{Hom}(X, -)$.

[1.0.1] **Claim:** The functor $\text{Hom}(X, -)$ is left exact. That is, a short exact sequence

$$0 \longrightarrow A \xrightarrow{i} B \xrightarrow{q} C \longrightarrow 0$$

gives an exact sequence

$$0 \longrightarrow \text{Hom}(X, A) \xrightarrow{i \circ -} \text{Hom}(X, B) \xrightarrow{q \circ -} \text{Hom}(X, C)$$

where the induced maps are by the obvious post-compositions with i and q . Similarly, the contravariant Hom functor $\text{Hom}(-, X)$ gives an exact sequence

$$0 \longrightarrow \text{Hom}(C, X) \xrightarrow{- \circ q} \text{Hom}(B, X) \xrightarrow{- \circ i} \text{Hom}(A, X)$$

where the induced maps are the pre-compositions with i and q .

Proof: For $f \in \text{Hom}(X, A)$, $i \circ f = 0$ implies $(i \circ f)(x) = 0$ for all $x \in X$, and then $f(x) = 0$ for all x since i is injective. Thus, $\text{Hom}(X, A) \rightarrow \text{Hom}(X, B)$ is injective, giving exactness at the left joint.

Since $q \circ i = 0$, any $f \in \text{Hom}(X, A)$ is mapped to $0 \in \text{Hom}(X, C)$ by $f \rightarrow q \circ i \circ f$. That is, the image of $i \circ -$ is contained in the kernel of $q \circ -$. On the other hand, for $g \in \text{Hom}(X, B)$ is mapped to $q \circ g = 0$ in $\text{Hom}(X, C)$,

$$g(X) \subset \ker q = \text{Im } i$$

Since i is injective, it is an isomorphism to its image, so there is an inverse $i^{-1} : \text{Im } i \rightarrow A$. Since $g(X) \subset \text{Im } i$, we can define

$$f = i^{-1} \circ g \in \text{Hom}(X, A)$$

[1] A functor $F : \mathcal{C} \rightarrow \mathcal{D}$ of categories whose hom-sets are *abelian groups additive* when the map on morphisms $\text{Hom}(A, B) \rightarrow \text{Hom}(FA, FB)$ given by F is a homomorphism of abelian groups. This also entails $F(A \oplus B) \approx FA \oplus FB$. These isomorphisms are required to be *natural*.

Certainly $i \circ f = g$, so the kernel is contained in the image. This gives exactness at the middle joint, and the left exactness. The exactness of the contravariant Hom is similar. ///

[1.0.2] Remark: The functor $\text{Hom}(X, -)$ is *not* right exact. For example, with

$$0 \longrightarrow \mathbb{Z} \xrightarrow{\times n} \mathbb{Z} \longrightarrow \mathbb{Z}/n \longrightarrow 0$$

with an integer $n > 1$, with $X = \mathbb{Z}/n$ there is no non-zero map of the torsion abelian group X to the free abelian group \mathbb{Z} . That is, the right joint in the following is not exact:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \text{Hom}(\mathbb{Z}/n, \mathbb{Z}) & \xrightarrow{\times n} & \text{Hom}(\mathbb{Z}/n, \mathbb{Z}) & \longrightarrow & \text{Hom}(\mathbb{Z}/n, \mathbb{Z}/n) \longrightarrow 0 \\
 & & \parallel & & \parallel & & \parallel \\
 & & 0 & & 0 & & \mathbb{Z}/n
 \end{array}$$

Similarly, the contravariant $\text{Hom}(-, X)$ is not right exact.

2. Adjointness and naturality

Two functors R and L (from \mathbb{Z} -modules to \mathbb{Z} -modules) are *mutually adjoint* when there is an *adjunction*, that is, an isomorphism of \mathbb{Z} -modules

$$\text{Hom}(LA, B) \approx \text{Hom}(A, RB) \quad (\text{for all } A, B)$$

The functor R is a *right adjoint*, and L is a *left adjoint*. The adjunction isomorphism is required to be *natural* or *functorial*, in the sense that, for each pair of morphisms $f : A' \rightarrow A$ and $g : B \rightarrow B'$ (yes, from A' to A , but from B to B') we have a commutative diagram [2]

$$\begin{array}{ccc}
 \text{Hom}(LA, B) & \xrightarrow{\approx} & \text{Hom}(A, RB) \\
 g \circ (-) \circ Lf \downarrow & & Rg \circ (-) \circ f \downarrow \\
 \text{Hom}(LA', B') & \xrightarrow{\approx} & \text{Hom}(A', RB')
 \end{array}$$

where the notation for pre-composition and post-composition is

$$g \circ (-) \circ Lf : F \longrightarrow g \circ F \circ Lf \quad (\text{for } F \in \text{Hom}(LA, B))$$

and

$$Rg \circ (-) \circ f : F \longrightarrow Rg \circ F \circ f \quad (\text{for } F \in \text{Hom}(A, RB))$$

In categories whose hom sets have additional structure, such as that of \mathbb{Z} -modules, the natural isomorphisms of hom sets are required to respect that additional structure, and satisfaction of this requirement is usually obvious.

We prove naturality in two examples of adjoint pairs.

[2.1] **Annihilated and co-annihilated modules** Let Λ be a ring with 1. Consider the category of Λ -modules and Λ -module homomorphisms. Fix an ideal I in Λ . A Λ -module N is *annihilated* by I if $i \cdot n = 0$ for all $i \in I$ and $n \in N$. For convenience, say that such N is *I-null*.

[2] Assembling these naturality isomorphisms into larger diagrams is critical in the later argument for left/right-exactness from adjointness.

Given a Λ -module M , M^I is a Λ -module with a Λ -module map $j : M^I \rightarrow M$ through which every map $N \rightarrow M$ from an I -null Λ -module N factors. Dually, M_I is an Λ -module with Λ -module map $q : M \rightarrow M_I$ through which every map $M \rightarrow N$ to an I -null Λ -module N factors.

Proof of *existence*, by construction, of M^I and M_I element-wise is easy, as sub-object and quotient, respectively:

$$\begin{cases} M^I &= \{m \in M : i \cdot m = 0 \text{ for all } i \in I\} \\ M_I &= M/(I \cdot M) \end{cases}$$

[2.1.1] **Claim:** The functors $LM = M_I$ and $RM = M^I$ are mutual adjoints:

$$\mathrm{Hom}_\Lambda(LA, B) \approx \mathrm{Hom}_\Lambda(A, RB)$$

Proof: The isomorphism of hom-sets is the obvious $f \rightarrow f \circ q$, where $q : A \rightarrow A_I$ is the quotient map. That $f \circ q$ has image inside the subobject $RB = B^I$ follows from the fact that $f : A \rightarrow B$ has image inside B^I , which follows from the fact that I acts by 0 on A_I .

The issue of interest is the naturality. Let $\alpha : A' \rightarrow A$ and $\beta : B \rightarrow B'$ be Λ -module homomorphisms. Let $q' : A' \rightarrow A_I$ be the quotient map. That $A \rightarrow A_I$ is a *functor* implicitly claims that there is a homomorphism $\alpha_I : A_I \rightarrow A_I$. Indeed, the composite

$$A' \rightarrow A \rightarrow A_I$$

must factor through A_I , by its universal property, yielding a unique $\alpha_I : A_I \rightarrow A_I$ fitting into the commutative diagram

$$\begin{array}{ccc} A' & \xrightarrow{q'} & A_I \\ \alpha \downarrow & & \downarrow \alpha_I \\ A & \xrightarrow{q} & A_I \end{array}$$

Similarly, there is $\beta^I : B^I \rightarrow B'^I$ fitting into a commutative diagram, namely, β^I is the restriction of β to B^I :

$$\begin{array}{ccc} B^I & \xrightarrow{i} & B \\ \beta^I \downarrow & & \downarrow \beta \\ B'^I & \xrightarrow{i'} & B' \end{array}$$

The *naturality* is the commutativity of

$$\begin{array}{ccc} \mathrm{Hom}(A_I, B) & \xrightarrow{\approx} & \mathrm{Hom}(A, B^I) \\ \beta \circ (-) \circ \alpha_I \downarrow & & \downarrow \beta^I \circ (-) \circ \alpha \\ \mathrm{Hom}(A_I, B') & \xrightarrow{\approx} & \mathrm{Hom}(A', B'^I) \end{array}$$

Note that at this point we do not give the top and bottom edge isomorphisms explicitly. This is clarified in the following.

The desired commutative diagram expands to a larger diagram upon making explicit the isomorphism whose

naturality is at issue. Namely, we claim that the following is commutative:

$$\begin{array}{ccccc}
 \mathrm{Hom}(A_I, B) & \xleftarrow[\approx]{i \circ -} & \mathrm{Hom}(A_I, B^I) & \xrightarrow[\approx]{- \circ q} & \mathrm{Hom}(A, B^I) \\
 \beta \circ - \downarrow & & \beta^I \circ - \downarrow & & \beta^I \circ - \downarrow \\
 \mathrm{Hom}(A_I, B') & \xleftarrow[\approx]{i \circ -} & \mathrm{Hom}(A_I, B'^I) & \xrightarrow[\approx]{- \circ q} & \mathrm{Hom}(A, B'^I) \\
 - \circ \alpha_I \downarrow & & - \circ \alpha_I \downarrow & & - \circ \alpha \downarrow \\
 \mathrm{Hom}(A'_I, B') & \xleftarrow[\approx]{i' \circ -} & \mathrm{Hom}(A'_I, B'^I) & \xrightarrow[\approx]{- \circ q'} & \mathrm{Hom}(A', B'^I)
 \end{array}$$

The three horizontal maps on the left half are the definition of $(-)^I$, while the three horizontal maps on the right half are the definition of $(-)_I$. Each vertical map is the image of a morphism by a *Hom* functor.

Commutativity of the upper left square follows from applying $\mathrm{Hom}(A_I, -)$ to the square expressing functoriality of $(-)_I$. Similarly, commutativity of the lower right square follows from applying $\mathrm{Hom}(-, B'^I)$ to the square expressing functoriality of $(-)^I$.

The upper right and lower left squares commute by associativity of composition of homs.

The directions of the horizontal maps in each row are in opposite directions, so they can be composed only because they are *isomorphisms*. The composition along the top edge, and the composition along the bottom edge, are the isomorphisms in the assertion of adjunction. The adjunction square is obtained by keeping only the four outer corner *Homs*, and the composite maps along the outer edges. ///

[2.2] Cartan-Eilenberg adjunction: $(-) \otimes X$ and $\mathrm{Hom}(X, -)$

[2.2.1] **Claim:** For all \mathbb{Z} -modules A, X, B there is a *natural* isomorphism of \mathbb{Z} -modules

$$\mathrm{Hom}(A \otimes X, B) \approx \mathrm{Hom}(A, \mathrm{Hom}(X, B))$$

Proof: Once one knows existence, there is only one possibility that makes sense: given $\Phi \in \mathrm{Hom}(A \otimes X, B)$, define $\varphi_\Phi \in \mathrm{Hom}(A, \mathrm{Hom}(X, B))$ by

$$\varphi_\Phi(a)(x) = \Phi(a \otimes x)$$

Conversely, given $\varphi \in \mathrm{Hom}(A, \mathrm{Hom}(X, B))$, there is $\Phi_\varphi \in \mathrm{Hom}(A \otimes X, B)$ by

$$\Phi_\varphi(a \otimes x) = \varphi(a)(x)$$

and extend by linearity. The maps $\Phi \rightarrow \varphi_\Phi$ and $\varphi \rightarrow \Phi_\varphi$ are mutual inverses. *Naturality* of the isomorphism $\varphi \rightarrow \Phi_\varphi$ and $\Phi \rightarrow \varphi_\Phi$ asserts the commutativity of diagrams attached to $f : A' \rightarrow A$ and $g : X' \rightarrow X$: there is a commutative diagram

$$\begin{array}{ccc}
 \mathrm{Hom}(A \otimes X, B) & \xrightarrow{\approx} & \mathrm{Hom}(A, \mathrm{Hom}(X, B)) \\
 h \circ (-) \circ (f \otimes g) \downarrow & & \downarrow a' \rightarrow (x' \rightarrow h((-)(fa'))(gx')) \\
 \mathrm{Hom}(A' \otimes X, B') & \xrightarrow{\approx} & \mathrm{Hom}(A', \mathrm{Hom}(X', B'))
 \end{array}$$

Note the awkwardness on the right-hand side of the diagram, leading to a more verbose description.

This is easy to check: starting with Φ in the upper left, going down gives $\Phi' = h \circ \Phi \circ (f \otimes g)$, and then going to the right gives φ' such that

$$\varphi'(a')(x') = \Phi'(a' \otimes x') = (h \circ \Phi \circ (f \otimes g))(a' \otimes x') = h\Phi(fa' \otimes gx')$$

Going the other way around the diagram, first we obtain φ such that $\varphi(a)(x) = \Phi(a \otimes x)$. Going down the right side gives φ' such that

$$\varphi'(a')(x') = \varphi(fa')(gx') = h\Phi(fa' \otimes gx')$$

The two outcomes are the same, which is the naturality. ///

3. A small Yoneda lemma

The innocent-seeming property of $\text{Hom}(X, -)$ below is a special case of *Yoneda's Lemma*. [3]

[3.0.1] **Theorem:** We have *sufficient* criteria for exactness: given A, B, C ,

$$\text{Hom}(X, A) \xrightarrow{f \circ -} \text{Hom}(X, B) \xrightarrow{g \circ -} \text{Hom}(X, C) \text{ exact for all } X \implies A \xrightarrow{f} B \xrightarrow{g} C \text{ exact}$$

Similarly,

$$\text{Hom}(C, X) \xrightarrow{- \circ g} \text{Hom}(B, X) \xrightarrow{- \circ f} \text{Hom}(A, X) \text{ exact for all } X \implies A \xrightarrow{f} B \xrightarrow{g} C \text{ exact}$$

[3.0.2] **Remark:** Exactness of $A \rightarrow B \rightarrow C$ does *not* imply exactness of the Hom diagram for all X . This was visible in proving *left* exactness of $M \rightarrow \text{Hom}(M, X)$.

Proof: On one hand, with $X = A$ and $F : X \rightarrow A$ the identity, exactness of the Hom sequence implies

$$0 = g \circ f \circ F = g \circ f$$

so $\text{Im } f \subset \ker g$. On the other hand, with $X = \ker g$ and $F : X \rightarrow B$ the inclusion, exactness of the Hom sequence (with $g \circ F = 0$) gives $F' : X \rightarrow A$ such that $f \circ F' = F$. Then

$$\ker g = \text{Im } F = \text{Im}(f \circ F') \subset \text{Im } f$$

The two containments together give $\ker g = \text{Im } f$, giving the result for covariant Hom.

For contravariant Hom $M \rightarrow \text{Hom}(M, X)$, with $X = C$ and $F : C \rightarrow X$ the identity, the exactness of the Hom sequence gives

$$0 = F \circ g \circ f = g \circ f$$

Thus, $\text{Im } f \subset \ker g$. On the other hand, with $X = B/\text{Im } f$ and $F : B \rightarrow X$ the quotient map, exactness of the Hom sequence gives $F' : C \rightarrow X$ such that $F' \circ g = F$. Thus, the kernel of g cannot be larger than $\text{Im } f$, or $F : B \rightarrow B/\text{Im } f$ could not factor through it. Thus, we have exactness. ///

[3] The functor $X \rightarrow \text{Hom}(X, -)$ from a category whose hom sets $\text{Hom}(A, B)$ are abelian groups, to the category of abelian groups, is an instance of a *Yoneda imbedding*.

4. Half-exactness of adjoints

A functor $F : \mathcal{C} \rightarrow \mathcal{D}$ on categories of modules is *additive*^[4] when

$$F(A \oplus B) \approx FA \oplus FB \quad (\text{for all } A, B \in \mathcal{C})$$

The isomorphism is required to be *natural*.

An *additive* functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is *left-exact* when^[5]

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0 \quad \text{exact} \quad \implies \quad 0 \rightarrow FA \rightarrow FB \rightarrow FC \quad \text{exact}$$

and *right-exact* when

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0 \quad \text{exact} \quad \implies \quad FA \rightarrow FB \rightarrow FC \rightarrow 0 \quad \text{exact}$$

[4.0.1] Theorem: Let L, R be mutually adjoint *additive* functors, with L the left and R the right adjoint. Then L is right half-exact and R is left half-exact. That is,

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0 \quad \text{exact} \quad \implies \quad LA \rightarrow LB \rightarrow LC \rightarrow 0 \quad \text{exact}$$

and

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0 \quad \text{exact} \quad \implies \quad 0 \rightarrow RA \rightarrow RB \rightarrow RC \quad \text{exact}$$

Proof: Left exactness of $M \rightarrow \text{Hom}(X, M)$ for any X applies to X replaced by LX , so

$$0 \rightarrow \text{Hom}(LX, A) \rightarrow \text{Hom}(LX, B) \rightarrow \text{Hom}(LX, C) \quad (\text{exact})$$

Adjointness of L and R , and *naturality* of the adjunction isomorphisms, give a commutative diagram with exact top row,

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Hom}(LX, A) & \longrightarrow & \text{Hom}(LX, B) & \longrightarrow & \text{Hom}(LX, C) \\ & & \downarrow \approx & & \downarrow \approx & & \downarrow \approx \\ 0 & \longrightarrow & \text{Hom}(X, RA) & \longrightarrow & \text{Hom}(X, RB) & \longrightarrow & \text{Hom}(X, RC) \end{array}$$

Then the bottom row is exact, for all X . By the small Yoneda lemma,

$$0 \rightarrow RA \rightarrow RB \rightarrow RC \quad (\text{exact})$$

^[4] More abstractly, a common structure on a category \mathcal{C} is that it be a *pre-additive* category, meaning that for $A, B \in \mathcal{C}$ the hom set $\text{Hom}_{\mathcal{C}}(A, B)$ is an *abelian group*, and the composition

$$\text{Hom}_{\mathcal{C}}(A, B) \times \text{Hom}_{\mathcal{C}}(B, C) \rightarrow \text{Hom}_{\mathcal{C}}(A, C)$$

is *bilinear*. Thus, more generally, for pre-additive categories \mathcal{C}, \mathcal{D} , a functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is *additive* when it preserves coproducts. In contemporary use, a category is *additive* when it is pre-additive *and* has a zero object, has finite coproducts. The pre-additive structure causes finite coproducts to be *products*, as well.

^[5] This is for *covariant* functors.

Similarly, for the other Hom functor, when in a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Hom}(C, RX) & \longrightarrow & \text{Hom}(B, RX, B) & \longrightarrow & \text{Hom}(A, RX) \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \text{Hom}(LC, X) & \longrightarrow & \text{Hom}(LB, X) & \longrightarrow & \text{Hom}(LC, X) \end{array}$$

the *top* row is exact, then the bottom row is exact. When this holds for all X , by the small Yoneda lemma,

$$LA \longrightarrow LB \longrightarrow LC \longrightarrow 0 \quad (\text{exact})$$

noting that this second Hom functor $M \rightarrow \text{Hom}(M, X)$ is *contravariant*. ///

[4.0.2] Corollary: The natural (adjunction) isomorphism $\text{Hom}(A \otimes X, B) \approx \text{Hom}(A, \text{Hom}(X, B))$ yields the *right* exactness of $M \rightarrow M \otimes X$. ///
