

NOTE ON 1-CROSSING PARTITIONS

M. BERGERSON, A. MILLER, A. PLIML, V. REINER, P. SHEARER, D. STANTON,
AND N. SWITALA

ABSTRACT. It is shown that there are $\binom{2n-r-1}{n-r}$ noncrossing partitions of an n -set together with a distinguished block of size r , and $\binom{n}{k-1}\binom{n-r-1}{k-2}$ of these have k blocks, generalizing a result of Bóna on partitions with one crossing. Furthermore, when one evaluates natural q -analogues of these formulae for q an n^{th} root of unity of order d , one obtains the number of such objects having d -fold rotational symmetry.

Given a partition π of the set $[n] := \{1, 2, \dots, n\}$, a *crossing* in π is a quadruple of integers (a, b, c, d) with $1 \leq a < b < c < d \leq n$ for which a, c are together in a block, and b, d are together in a different block. It is well-known [10, Exercises 6.19(pp)], [4] that the number of *noncrossing partitions* of $[n]$ (that is, those with no crossings) is the Catalan number $C_n = \frac{1}{n+1}\binom{2n}{n}$, and the number of noncrossing partitions of $[n]$ into k blocks is the Narayana number $\frac{1}{n}\binom{n}{k-1}\binom{n}{k}$.

Our starting point is the more recent observation of Bóna [2, Theorem 1] that the number of partitions of $[n]$ having *exactly one* crossing has the even simpler formula $\binom{2n-5}{n-4}$. Bóna's proof utilizes the fact that C_n is also well-known to count triangulations of a convex $(n+2)$ -gon; this allows him to biject 1-crossing partitions of $[n]$ to dissections of an n -gon that use exactly $n-4$ diagonals. The proof is then completed by plugging $d = n-4$ into the formula $\frac{1}{d+1}\binom{n+d-1}{d}\binom{n-3}{d}$ of Kirkman (first proven by Cayley; see [7]) for the number of dissections of an n -gon using d diagonals.

The goal here is to generalize Bóna's result to count 1-crossing partitions by their number of blocks, and also to examine a natural q -analogue with regard to the *cyclic sieving phenomenon* shown in [8] for certain q -Catalan and q -Narayana numbers. The crux is the observation that 1-crossing partitions of $[n]$ biject naturally with noncrossing partitions of $[n]$ having a distinguished 4-element block: replace the crossing pair of blocks $\{a, c\}, \{b, d\}$ with a single distinguished block $\{a, b, c, d\}$. Thus one should count the following more general objects.

Definition 1. An *r -blocked noncrossing partition* of $[n]$ is a pair (π, B) of a noncrossing partition π together with a distinguished r -element block B of π .

Note that the notion of a crossing in a partition is invariant under cyclic rotations $i \mapsto i+1 \pmod n$ of the set $[n]$. Consequently the cyclic group $C = \mathbb{Z}_n$ acts on the

Date: August 2006.

Key words and phrases. noncrossing partition, cyclic sieving phenomenon.

This work was the result of an REU at the University of Minnesota School of Mathematics in Summer 2006, mentored by V. Reiner and D. Stanton, and supported by NSF grants DMS-0601010 and DMS-0503660. The authors also thank D. Armstrong for helpful conversations.

set of r -blocked noncrossing partition of $[n]$, preserving the number of blocks. Also define $\begin{bmatrix} n \\ k \end{bmatrix}_q := \frac{[n]_q!}{[k]_q! [n-k]_q!}$, where $[n]_q! := [n]_q [n-1]_q \cdots [2]_q [1]_q$ with $[n]_q := \frac{1-q^n}{1-q}$.

Theorem 1. *The number of r -blocked noncrossing partition of $[n]$, and the number with exactly k blocks, are given by the formulae*

$$(0.1) \quad a(n, r) := \binom{2n-r-1}{n-r}, \quad a(n, k, r) := \binom{n}{k-1} \binom{n-r-1}{k-2}.$$

Furthermore, for any d dividing n , the number of d -fold symmetric r -blocked noncrossing partitions of $[n]$, and the number having exactly k blocks, are obtained by plugging in any primitive d^{th} root-of-unity for q in the natural q -analogues

$$(0.2) \quad a_q(n, r) := \begin{bmatrix} 2n-r-1 \\ n-r \end{bmatrix}_q, \quad a_q(n, k, r) := q^{(k-1)(k-2)} \begin{bmatrix} n \\ k-1 \end{bmatrix}_q \begin{bmatrix} n-r-1 \\ k-2 \end{bmatrix}_q.$$

Note that taking $r = 4$ and replacing k by $k - 1$ in (0.1), one finds agreement with Bóna's count of $\binom{2n-5}{n-4}$, as well as the formula $\binom{n}{k-2} \binom{n-5}{k-3}$ for the number of 1-crossing partitions with k blocks.

Proof. (of Theorem 1) For the first assertion, it suffices to prove the formula for $a(n, k, r)$; the formula for $a(n, r)$ follows from the Chu-Vandermonde summation.

Let $A(n, k, r)$ be the set of r -blocked noncrossing partitions of $[n]$ with k blocks, which we wish to count. Let $B(n, k, r)$ be the set of triples (π, B, i) in which π is a noncrossing partition of $[n]$ with k blocks, i is a chosen element of $[n]$, and B is an r -element block of π , with $i \in B$. Let $C(n, k, r)$ be the set of noncrossing partitions of $[n]$ in which the element 1 lies in an r -element block.

Counting $B(n, k, r)$ in two ways, one finds

$$r \cdot |A(n, k, r)| = |B(n, k, r)| = n \cdot |C(n, k, r)|,$$

and hence $a(n, k, r) = |A(n, k, r)| = \frac{n}{r} |C(n, k, r)|$.

To count $|C(n, k, r)|$, note that Dershowitz and Zaks [4] give a bijection between noncrossing partitions and ordered trees, which restricts to a bijection between $C(n, k, r)$ and the set $D(n, k, r)$ of all ordered trees having n edges, root degree r , and k internal nodes. On the other hand, the set $D(n, k, r)$ has been enumerated multiple times in the literature via generating functions and Lagrange inversion (e.g. in [3, 5]), and can also be done semi-bijectively (see [1]):

$$|D(n, k, r)| = \frac{r}{n} \binom{n}{k-1} \binom{n-r-1}{k-2}.$$

Combining this with the foregoing proves (0.1).

For the assertion about q -analogues, we first deal with the case of $a_q(n, k, r)$. Note that for any d dividing n , an r -blocked noncrossing partition of $[n]$ having k blocks has no chance of being d -fold symmetric unless r is divisible by d and k is congruent to 1 mod d . Furthermore, when these congruences hold, if one defines $n' := \frac{n}{d}$, $r' := \frac{r}{d}$, $k' := \frac{k-1}{d}$, then the quotient mapping $[n] \cong \mathbb{Z}_n \rightarrow \mathbb{Z}_{n'} \cong [n']$ gives a natural bijection between d -fold symmetric r -blocked noncrossing partitions of $[n]$ with k blocks, and r' -blocked noncrossing partitions of $[n']$ with $k' + 1$ blocks. Hence by the first part of the theorem, there are exactly $\binom{n'}{k'} \binom{n'-r'-1}{k'-1}$ such d -fold symmetric r -blocked noncrossing partition of $[n]$ having k blocks in this case.

On the other hand, one can easily evaluate $a_q(n, k, r)$ when q is a primitive d^{th} root-of-unity for d dividing n , using the q -Lucas theorem (Lemma 2 below). One finds that it vanishes unless r is divisible by d and k is congruent to 1 mod d , in which case it equals $\binom{n'}{k'} \binom{n'-r'-1}{k'-1}$, as desired.

For the assertion about $a_q(n, r)$, one can either argue in a similar fashion, or use the identity $\left[\begin{matrix} 2n-r-1 \\ n-r \end{matrix} \right]_q = \sum_k q^{(k-1)(k-2)} \left[\begin{matrix} n \\ k-1 \end{matrix} \right]_q \left[\begin{matrix} n-r-1 \\ k-2 \end{matrix} \right]_q$, an instance of the q -Chu-Vandermonde summation; see e.g. [6, (7.6)]. \square

The following straightforward lemma used in the above proof has been rediscovered many times; see [9, Theorem 2.2] for a proof and some history.

Lemma 2. (q -Lucas theorem) *Given nonnegative integers n, k, d , with $1 \leq d \leq n$, uniquely write $n = n'd + n''$ and $k = k'd + k''$ with $0 \leq n'', k'' < d$. If q is a primitive d^{th} root-of-unity, then*

$$\left[\begin{matrix} n \\ k \end{matrix} \right]_q = \binom{n'}{k'} \left[\begin{matrix} n'' \\ k'' \end{matrix} \right]_q.$$

One can derive an explicit formula for the number of 2-crossing partitions of $[n]$, but it is much messier than $a(n, r)$ above, and appears to have no q -analogue with good behavior. However, Bóna [2] does show that for each fixed k , the generating function counting k -crossing partitions of $[n]$ is a rational function of x and $\sqrt{1-4x}$.

REFERENCES

- [1] M. Bergerson, A. Miller, P. Shearer and N. Switala, Enumeration of 1-crossing and 2-crossing partitions with refinements. REU report, Summer 2006. Available at www.math.umn.edu/reiner/REU/REU.html.
- [2] M. Bóna, Partitions with k Crossings. *The Ramanujan Journal* **3** (1999), 215-220.
- [3] H. Burgiel and V. Reiner, Two signed associahedra. *New York J. Math. (electronic)* **4** (1998), 83-95.
- [4] N. Dershowitz and S. Zaks, Ordered Trees and Non-Crossing Partitions. *Disc. Math.* **62** (1986), 215-218.
- [5] E. Deutsch, Dyck Path Enumeration. *Disc. Math.* **204** (1999), 167-202.
- [6] V. Kac and P. Cheung, *Quantum calculus*. Universitext. Springer-Verlag, New York, 2002.
- [7] J. Przytycki and A.S. Sikora, Adam S, Polygon dissections and Euler, Fuss, Kirkman, and Cayley numbers. *J. Combin. Theory Ser. A92* (2000), 68-76.
- [8] V. Reiner, D. Stanton, and D. White, The cyclic sieving phenomenon, *Journal of Combinatorial Theory, Series A* **108** (2004), 17-50.
- [9] B. Sagan, Congruence properties of q -analogs, *Adv. Math.* **95** (1992), 127-143.
- [10] R.P. Stanley, *Enumerative Combinatorics, Vol. 2*. Cambridge Studies in Advanced Mathematics **62**, Cambridge University Press, Cambridge, 1999.