

# Orthogonal basic hypergeometric Laurent polynomials

Mourad E.H. ISMAIL <sup>†</sup> and Dennis STANTON <sup>‡</sup>

<sup>†</sup> *Department of Mathematics, University of Central Florida, Orlando, FL 32816, USA, and  
Department of Mathematics, King Saud University, Riyadh, Saudi Arabia*

E-mail: [mourad.eh.ismail@gmail.com](mailto:mourad.eh.ismail@gmail.com)

<sup>‡</sup> *School of Mathematics, University of Minnesota, Minneapolis, MN 55455, USA*

E-mail: [stanton@math.umn.edu](mailto:stanton@math.umn.edu)

Received ???, in final form ???; Published online ???

doi:10.3842/SIGMA.201\*.\*\*\*

**Abstract.** The Askey-Wilson polynomials are orthogonal polynomials in  $x = \cos \theta$ , which are given as a terminating  ${}_4\phi_3$  basic hypergeometric series. The non-symmetric Askey-Wilson polynomials are Laurent polynomials in  $z = e^{i\theta}$ , which are given as a sum of two terminating  ${}_4\phi_3$ 's. They satisfy a biorthogonality relation. In this paper new orthogonality relations for single  ${}_4\phi_3$ 's which are Laurent polynomials in  $z$  are given, which imply the non-symmetric Askey-Wilson biorthogonality. These results include discrete orthogonality relations. They can be considered as a classical analytic study of the results for non-symmetric Askey-Wilson polynomials which were previously obtained by affine Hecke algebra techniques.

*Key words:* Askey-Wilson polynomials; orthogonality

*2010 Mathematics Subject Classification:* 33D45

## 1 Introduction

The Askey-Wilson polynomials  $p_n(x; \mathbf{t} | q)$  are polynomials in  $x = \cos \theta$ , and depend upon parameters  $q$ ,  $t_1$ ,  $t_2$ ,  $t_3$ , and  $t_4$ . They may be defined by the terminating basic hypergeometric series [1], [8, §15.2]

$$p_n(x; \mathbf{t} | q) = t_1^{-n} (t_1 t_2, t_1 t_3, t_1 t_4; q)_n \times {}_4\phi_3 \left( \begin{matrix} q^{-n}, t_1 t_2 t_3 t_4 q^{n-1}, t_1 e^{i\theta}, t_1 e^{-i\theta} \\ t_1 t_2, t_1 t_3, t_1 t_4 \end{matrix} \middle| q, q \right). \quad (1.1)$$

They are orthogonal polynomials in  $x$ , and have a known orthogonality relation (see (2.1)).

Clearly the Askey-Wilson polynomials are also Laurent polynomials in  $z = e^{i\theta}$ , since  $2x = z + 1/z$ . One may ask if there is a natural basis for Laurent polynomials in  $z$  which also satisfy orthogonality relations. The non-symmetric Askey-Wilson polynomials [16],[17] are Laurent polynomials, which are biorthogonal with respect to a modified Askey-Wilson weight, are one answer to this question. They may be explicitly given as a sum of two terminating basic hypergeometric series.

The purpose of this paper is to use classical analytic methods to study the orthogonality relations for single  ${}_4\phi_3$ 's which imply the non-symmetric Askey-Wilson polynomial biorthogonality. The main orthogonality results are Theorem 1, Theorem 2, Theorem 3, and Theorem 4. We show that special function techniques, explicit summations and self-adjoint operators, may be applied to derive these orthogonalities. We explicitly find three and four term relations for Laurent polynomials in §4. As a corollary of our computations we rederive the  $L^2$ -norm of the non-symmetric Askey-Wilson polynomials in §6, and reprove the creation type formulas for the Laurent polynomials in §5. Asymptotics are given in Theorem 10, and Racah orthogonality

in Theorem 11. This work may be considered as a classical study of the one variable case in Cherednik [6], Macdonald [16], carefully studied by Noumi and Stokman [17], who considered many of these results from the point of view of affine Hecke algebras.

There are different theories of orthogonal polynomials, biorthogonal polynomials and rational functions which originate from different types of continued fractions. The  $J$ -fractions lead to orthogonal polynomials on the line while the  $PC$  fractions lead to orthogonal polynomials on the unit circle, [11], [14]–[15], [8]. The theory of  $T$ -fractions leads to orthogonal Laurent polynomials and is explained in the book [14]–[15], and the excellent survey article [5]. Ismail and Masson, [9] introduced the more general  $R_I$  and  $R_{II}$  fractions which naturally lead to biorthogonal orthogonal rational functions. A. Zhedanov [21] pointed out the connection of the latter continued fractions and the generalized eigenvalue problem [19]. For applications of the  $R_I$  and  $R_{II}$  fractions to integrable systems, see [18]. There is a more abstract approach to biorthogonality presented in Brezinski's monograph [4] with many applications to numerical analysis and computational mathematics.

## 2 Notation

Let  $V_n$  be the  $(2n + 1)$ -dimensional real vector space spanned by the Laurent polynomials  $\{z^{-n}, z^{1-n}, \dots, z^{n-1}, z^n\}$ . Most of this section is devoted to notation for specific spanning sets in  $V_n$ . These sets have orthogonality relations given in later sections, we define the appropriate measures here.

We assume throughout the paper that  $x = \cos \theta$  and  $z = e^{i\theta}$ , and  $\mathbf{t}$  stands for the vector  $(t_1, t_2, t_3, t_4)$ . We shall always assume that  $0 < q < 1$  and each  $t_i$  is real with  $|t_i| < 1$ .

The weight function for the Askey-Wilson polynomials is given by

$$w_{\text{aw}}(x; \mathbf{t}|q) = \frac{(q, e^{2i\theta}, e^{-2i\theta}; q)_\infty}{\prod_{j=1}^4 (t_j e^{i\theta}, t_j e^{-i\theta}; q)_\infty}.$$

The Askey-Wilson orthogonality [8, (15.2.4)] is

$$\begin{aligned} \frac{1}{2\pi} \int_{-1}^1 p_m(x; \mathbf{t}|q) p_n(x; \mathbf{t}|q) w_{\text{aw}}(x; \mathbf{t}|q) \frac{dx}{\sqrt{1-x^2}} = \\ \prod_{1 \leq j < k \leq 4} (t_j t_k; q)_n \frac{(q, t_1 t_2 t_3 t_4 q^{n-1}; q)_n}{(t_1 t_2 t_3 t_4; q)_{2n}} \mu(\mathbf{t}|q) \delta_{m,n}, \end{aligned} \quad (2.1)$$

while the Askey-Wilson integral evaluation is [3], [8]

$$\frac{1}{2\pi} \int_0^\pi \frac{(q, e^{2i\theta}, e^{-2i\theta}; q)_\infty}{\prod_{j=1}^4 (t_j e^{i\theta}, t_j e^{-i\theta}; q)_\infty} d\theta = \frac{(t_1 t_2 t_3 t_4; q)_\infty}{\prod_{1 \leq j < k \leq 4} (t_j t_k; q)_\infty} = \mu(\mathbf{t}|q). \quad (2.2)$$

The orthogonality relations for Laurent polynomials in  $V_n$  uses a small variation of the Askey-Wilson weight, which is given by

$$\begin{aligned} w_{\text{cher}}(z; \mathbf{t}|q) &= w_{\text{aw}}(x; \mathbf{t}|q) \frac{(1 - t_1/z)(1 - t_2/z)}{1 - z^{-2}} \\ &= \frac{(q, z^2, qz^{-2}; q)_\infty}{\prod_{j=1}^4 (t_j z; q)_\infty \prod_{k=3}^4 (t_k/z, qt_{k-2}/z; q)_\infty}. \end{aligned}$$

Even though this weight is not positive on the unit circle, and does not lead to a positive definite quadratic form, we will define a bilinear form using  $w_{\text{cher}}(z; \mathbf{t}|q)$ , and find orthogonal bases for  $V_n$  in §3 and §4.

The orthogonality relations will involve Laurent polynomials in  $z$  which may be defined in terms of  ${}_4\phi_3$  functions. We shall need four of these functions:  $R_n, S_n, T_n$ , and  $U_n$ .

The first polynomial  $R_n(z, \mathbf{t}|q)$  is an unscaled version of the Askey-Wilson polynomials

$$R_n(z; \mathbf{t}|q) = {}_4\phi_3 \left( \begin{matrix} q^{-n}, t_1 t_2 t_3 t_4 q^{n-1}, t_1 z, t_1/z \\ t_1 t_2, t_1 t_3, t_1 t_4 \end{matrix} \middle| q, q \right). \quad (2.3)$$

Put  $S_0(z) = 0$ , and for  $n \geq 1$ ,

$$S_n(z; \mathbf{t}|q) = z(1 - t_3/z)(1 - t_4/z) {}_4\phi_3 \left( \begin{matrix} q^{1-n}, t_1 t_2 t_3 t_4 q^n, t_1 z, qt_1/z \\ qt_1 t_2, qt_1 t_3, qt_1 t_4 \end{matrix} \middle| q, q \right), \quad (2.4)$$

$$T_n(z; \mathbf{t}|q) = {}_4\phi_3 \left( \begin{matrix} q^{-n}, t_1 t_2 t_3 t_4 q^{n-1}, t_1 z, qt_1/z \\ qt_1 t_2, t_1 t_3, t_1 t_4 \end{matrix} \middle| q, q \right), \quad (2.5)$$

and  $U_0(z) = 0$ , and for  $n \geq 1$ ,

$$U_n(z; \mathbf{t}|q) = \frac{1}{z}(1 - t_1 z)(1 - t_2 z) R_{n-1}(z; qt_1, qt_2, t_3, t_4|q). \quad (2.6)$$

The above representation (2.6) expresses  $U_n$  as a multiple of a scaled  $R_n$ . The  $S_n$  and  $T_n$  in (2.4) and (2.5) can also be written in terms of scaled  $R_n$ 's as follows

$$\begin{aligned} S_n(z; \mathbf{t}|q) &= z(1 - t_3/z)(1 - t_4/z) T_{n-1}(z; t_1, t_2, qt_3, qt_4), \\ T_n(z; \mathbf{t}|q) &= R_n(q^{-1/2}z; q^{1/2}t_1, q^{1/2}t_2, q^{-1/2}t_3, q^{-1/2}t_4). \end{aligned}$$

In other words

$$S_n(z; \mathbf{t}|q) = z(1 - t_3/z)(1 - t_4/z) R_{n-1}(q^{-1/2}z; q^{1/2}t_1, q^{1/2}t_2, q^{1/2}t_3, q^{1/2}t_4).$$

Clearly

$$\{R_0, \dots, R_n, S_1, \dots, S_n, T_0, \dots, T_n, U_1, \dots, U_n\} \subset V_n,$$

but this set could not form a basis for  $V_n$  for  $n \geq 0$ . There are very specific dependencies amongst these Laurent polynomials which follow from contiguous relations. It follows from Proposition 1 that for  $n \geq 1$  the subspace  $W_n = \text{span}\{R_n, S_n, T_n, U_n\}$  of  $V_n$  is 2-dimensional. Any two of these four Laurent polynomials are linearly independent.

We need some contiguous relations for balanced  ${}_4\phi_3$ 's. Let

$$\phi(a, b, c, d; e, f, g) = {}_4\phi_3 \left( \begin{matrix} a, b, c, d \\ e, f, g \end{matrix} \middle| q, q \right)$$

denote a  ${}_4\phi_3$  which is terminating and balanced. By  $\phi_+$  we mean change  $a, b, \dots, g$  by  $aq, bq, \dots, gq$ , respectively, so  $\phi_+$  is no longer balanced. When a parameter  $\alpha$  is changed to  $\alpha q^{\pm 1}$  we write  $\phi(\alpha \pm)$ . Wilson [20] proved contiguous relations which imply

$$\phi(a+, e+) - \phi = \frac{q(a-e)(1-b)(1-c)(1-d)}{(1-e)(1-eq)(1-f)(1-g)} \phi_+(e+), \quad (2.7)$$

$$\begin{aligned} \frac{a(1-f/a)(1-g/a)}{(1-f)(1-g)} \phi_+(a-) + \frac{(b-e)(1-c/e)}{(1-b)(1-c)} \phi(d+, e+) \\ = \frac{(1-e)(1-bc/e)}{(1-b)(1-c)} \phi. \end{aligned} \quad (2.8)$$

**Proposition 1.** *The following connection relations hold for  $n \geq 1$ ,*

$$\begin{aligned} T_n(z; \mathbf{t}|q) - R_n(z; \mathbf{t}|q) &= \frac{qt_1(1 - q^{-n})(1 - t_1t_2t_3t_4q^{n-1})}{(1 - t_1t_2)(1 - qt_1t_2)(1 - t_1t_3)(1 - t_1t_4)} U_n(z; \mathbf{t}|q), \\ (1 - t_1t_3)(1 - t_1t_4)R_n(z; \mathbf{t}|q) - t_1S_n(z; \mathbf{t}|q) &= \frac{t_1(1 - t_1t_2q^n)(1 - t_3t_4q^{n-1})}{q^{n-1}(1 - t_1t_2)(1 - qt_1t_2)} U_n(z; \mathbf{t}|q), \\ \frac{t_1}{(1 - t_1t_3)(1 - t_1t_4)} S_n(z; \mathbf{t}|q) - \frac{(1 - q^n t_1 t_2)(1 - q^{n-1} t_3 t_4)}{(1 - q^n)(1 - t_1 t_2 t_3 t_4 q^{n-1})} T_n(z; \mathbf{t}|q) \\ &+ \frac{q^n(1 - t_1 t_2)(1 - t_3 t_4 / q)}{(1 - q^n)(1 - t_1 t_2 t_3 t_4 q^{n-1})} R_n(z; \mathbf{t}|q) = 0. \end{aligned}$$

**Proof.** The first statement follows from (2.7) with  $a = t_1/z$ ,  $b = q^{-n}$ ,  $c = t_1t_2t_3t_4q^{n-1}$ ,  $d = t_1z$ ,  $e = t_1t_2$ ,  $f = t_1t_3$ , and  $g = t_1t_4$ . The third statement follows from (2.8) with  $a = t_1z$ ,  $b = q^{-n}$ ,  $c = t_1t_2t_3t_4q^{n-1}$ ,  $d = t_1/z$ ,  $e = t_1t_2$ ,  $f = t_1t_3$ , and  $g = t_1t_4$ . The second statement follows from the other two.  $\blacksquare$

We will also use the Sears transformation [7, (III.15)],

$$\begin{aligned} & {}_4\phi_3 \left( \begin{matrix} q^{-n}, A, B, C \\ D, E, F \end{matrix} \middle| q, q \right) \\ &= \frac{(E/A, F/A; q)_n}{(E, F; q)_n} A^n {}_4\phi_3 \left( \begin{matrix} q^{-n}, A, D/B, D/C \\ D, Aq^{1-n}/E, Aq^{1-n}/F \end{matrix} \middle| q, q \right), \end{aligned} \quad (2.9)$$

where  $DEF = ABCq^{1-n}$ . The Sears transformation (2.9) applied to the  $R_n$  and  $T_n$  functions give

$$R_n(z; t_1, t_2, t_3, t_4|q) = \frac{t_1^n(t_2t_3, t_3t_4; q)_n}{t_3^n(t_1t_2, t_1t_4; q)_n} R_n(z; t_3, t_2, t_1, t_4|q) \quad (2.10)$$

and

$$T_n(z; t_1, t_2, t_3, t_4|q) = \frac{t_1^n(t_2t_3, t_2t_4; q)_n}{t_2^n(t_1t_3, t_1t_4; q)_n} T_n(z; t_2, t_1, t_3, t_4|q). \quad (2.11)$$

One may also easily see from the definition of  $R_n$  and  $S_n$  that

$$\begin{aligned} R_n(1/z; 1/\mathbf{t}|1/q) &= R_n(z; \mathbf{t}|q), \\ S_n(1/z; 1/\mathbf{t}|1/q) &= S_n(z; \mathbf{t}|q)/t_3t_4. \end{aligned} \quad (2.12)$$

### 3 Orthogonality Relations

The main result of this section is Theorem 1.

For Laurent polynomials  $f(z)$  and  $g(z)$  define a bilinear form by

$$\langle f, g \rangle_{\text{cher}} = \frac{1}{2\pi i} \oint_{|z|=1} f(z) g(z) w_{\text{cher}}(z; \mathbf{t}|q) \frac{dz}{z}.$$

**Theorem 1.** *The set  $\{R_0, \dots, R_n, U_1, \dots, U_n\}$  is an orthogonal basis for  $V_n$  with respect to the symmetric bilinear form  $\langle, \rangle_{\text{cher}}$ . Moreover,  $\langle R_m, R_m \rangle_{\text{cher}}$  and  $\langle U_m, U_m \rangle_{\text{cher}}$  are non-zero.*

The proof of Theorem 1 will be accomplished in three stages. We get  $R_n - R_m$  orthogonality for “free”. We establish in Propositions 2 and 3 orthogonalities between  $R_n$  and  $T_m$ , which changes into  $R_n - U_m$  and  $U_n - U_m$  orthogonality.

First we establish  $R_n - R_m$  orthogonality from Askey-Wilson orthogonality. It is a restatement of (2.1), which was given in [17, Lemma 6.4].

**Proposition 2.** *The orthogonality relation for the  $R_n$  functions is*

$$\langle R_m, R_n \rangle_{\text{cher}} = \frac{(1 - t_1 t_2) t_1^{2n} (q, t_2 t_3, t_2 t_4, t_3 t_4, t_1 t_2 t_3 t_4 q^{n-1}; q)_n \mu(\mathbf{t}|q) \delta_{m,n}}{(t_1 t_2, t_1 t_3, t_1 t_4; q)_n (t_1 t_2 t_3 t_4; q)_{2n}}$$

**Proof.** Rewrite the left-hand side of Proposition 2 as

$$\begin{aligned} & \frac{1}{2\pi} \int_{-\pi}^{\pi} R_m(e^{i\theta}; \mathbf{t}|q) R_n(e^{i\theta}; \mathbf{t}|q) w_{\text{aw}}(\cos \theta; \mathbf{t}|q) \\ & \quad \times \frac{e^{i\theta} - (t_1 + t_2) + t_1 t_2 e^{-i\theta}}{2 \sin \theta} d\theta \\ &= \frac{1 - t_1 t_2}{2\pi} \int_0^{\pi} R_m(e^{i\theta}; \mathbf{t}|q) R_n(e^{i\theta}; \mathbf{t}|q) w_{\text{aw}}(\cos \theta; \mathbf{t}|q) d\theta, \end{aligned}$$

since  $R_n(e^{i\theta}; \mathbf{t}|q)$  and  $w_{\text{aw}}$  are even functions of  $\theta$ . ■

To establish the  $U_n - R_m$  orthogonality we first give an integral evaluation for “gluing” a specific Laurent polynomial onto the measure.

**Lemma 1.** *For non-negative integers  $k$  and  $j$  we have*

$$\begin{aligned} & \frac{1}{2\pi i} \oint_{|z|=1} (t_1 z, q t_1/z; q)_j (t_3 z, t_3/z; q)_k w_{\text{cher}}(z; \mathbf{t}|q) \frac{dz}{z} = \\ & \frac{(t_1 t_2 t_3 t_4 q^{j+k}; q)_{\infty}}{(t_1 t_2 q^{j+1}, t_1 t_3 q^{j+k}, t_1 t_4 q^j, t_2 t_3 q^k, t_2 t_4, t_3 t_4 q^k; q)_{\infty}}. \end{aligned}$$

**Proof.** The integral is

$$\begin{aligned} & \frac{1}{2\pi i} \oint_{|z|=1} \frac{(q, z^2, z^{-2}; q)_{\infty}}{(t_1 q^j z, t_2 z, t_3 q^k z, t_4 z, q^j t_1/z, t_2/z, q^k t_3/z, t_4/z; q)_{\infty}} \\ & \quad \times \frac{(1 - t_2/z)(1 - q^j t_1/z)}{(1 - z^{-2})} \frac{dz}{z}. \end{aligned}$$

This time the rational function in  $z = e^{i\theta}$  in the integrand is

$$\frac{e^{i\theta} - (q^j t_1 + t_2) + t_1 t_2 q^j e^{-i\theta}}{2i \sin(\theta)}. \quad (3.1)$$

The remaining part of the integrand is symmetric in  $\theta$ , so only the odd part of the numerator in (3.1) contributes. The Askey-Wilson integral evaluation gives the result.

$$\begin{aligned} &= \frac{1}{2} (1 - t_1 t_2 q^j) \frac{1}{2\pi} \int_{-\pi}^{\pi} w_{\text{aw}}(\cos \theta; q^j t_1, t_2, q^k t_3, t_4|q) d\theta \\ &= \frac{(t_1 t_2 t_3 t_4 q^{j+k}; q)_{\infty}}{(t_1 t_2 q^{j+1}, t_1 t_3 q^{j+k}, t_1 t_4 q^j, t_2 t_3 q^k, t_2 t_4, t_3 t_4 q^k; q)_{\infty}}. \end{aligned}$$
■

**Proposition 3.** *We have the orthogonality relation*

$$\langle R_m, T_n \rangle_{\text{cher}} = \langle R_m, R_n \rangle_{\text{cher}} \delta_{m,n}.$$

**Proof.** The proof is done in three stages:  $m < n$ ,  $m = n$ , and  $m > n$ .

Let  $I_{j,k}$  be the integral in Lemma 1. Using the explicit definition of  $T_n(z; \mathbf{t} | q)$  we have

$$\begin{aligned} & \frac{1}{2\pi i} \oint_{|z|=1} (t_3 z, t_3/z; q)_k T_n(z; \mathbf{t} | q) w_{\text{cher}}(z; \mathbf{t} | q) \frac{dz}{z} \\ &= \sum_{j=0}^n \frac{(q^{-n}, t_1 t_2 t_3 t_4 q^{n-1}; q)_j}{(q, q t_1 t_2, t_1 t_3, t_1 t_4; q)_j} q^j I_{j,k} \\ &= \frac{(t_1 t_2 t_3 t_4 q^k; q)_\infty}{(q, q t_1 t_2, q^k t_1 t_3, t_1 t_4, q^k t_2 t_3, t_2 t_4, q^k t_3 t_4; q)_\infty} {}_3\phi_2 \left( \begin{matrix} q^{-n}, t_1 t_2 t_3 t_4 q^{n-1}, q^k t_1 t_3 \\ t_1 t_3, t_1 t_2 t_3 t_4 q^k \end{matrix} \middle| q, q \right). \end{aligned}$$

From the  $q$ -Pfaff-Saalschütz theorem [7, (II.12)], the sum of the above  ${}_3\phi_2$  contains a factor  $(q^{-k}; q)_n$  which vanishes for  $k < n$ . Thus if  $m < n$ , then  $\langle R_m, T_n \rangle_{\text{cher}} = 0$ .

When  $k = n$  the the  $q$ -Pfaff-Saalschütz theorem [7, (II.12)] evaluates the above integral to

$$\begin{aligned} & \frac{1}{2\pi i} \oint_{|z|=1} (t_3 z, t_3/z; q)_n T_n(z; \mathbf{t} | q) w_{\text{cher}}(z; \mathbf{t} | q) \frac{dz}{z} \\ &= \frac{(t_1 t_2 t_3 t_4 q^{2n}; q)_\infty (-t_1 t_3)^n q^{\binom{n}{2}}}{(q^{n+1}, q t_1 t_2, t_1 t_3, t_1 t_4, t_2 t_3 q^n, t_2 t_4 q^n, t_3 t_4 q^n; q)_\infty}. \end{aligned}$$

Using (2.10) an explicit computation shows that Proposition 3 holds for  $m = n$ .

Next we consider the case  $m > n$ . In this case consider the integral

$$I = \frac{1}{2\pi i} \oint_{|z|=1} (t_1 z, q t_1/z; q)_j R_m(z; \mathbf{t} | q) w_{\text{cher}}(z; \mathbf{t} | q) \frac{dz}{z}$$

After using (2.10) for  $R_m$ ,  $I$  is a constant multiple of

$$\sum_{k=0}^m \frac{(q^{-m}, t_1 t_2 t_3 t_4 q^{m-1}; q)_k q^k}{(q, t_1 t_3, t_2 t_3, t_3 t_4; q)_k} I_{j,k}.$$

Applying Lemma 1 we see that the integral  $I$  is a constant multiple of

$${}_3\phi_2 \left( \begin{matrix} q^{-m}, t_1 t_2 t_3 t_4 q^{m-1}, q^j t_1 t_3 \\ t_1 t_3, t_1 t_2 t_3 t_4 q^j \end{matrix} \middle| q, q \right),$$

which again by the  $q$ -Pfaff-Saalschütz theorem [7, (II.12)] produces a factor  $(q^{-j}; q)_m$  and therefore vanishes for  $j < m$ .  $\blacksquare$

Finally we come to the  $T_n - T_m$  orthogonality, and finding the  $L^2$ -norms.

**Proposition 4.** *We have the orthogonality relation*

$$\langle T_m, T_n \rangle_{\text{cher}} = \frac{(1 - t_1 t_2)(1 - t_3 t_4/q)}{(1 - t_1 t_2 q^n)(1 - t_3 t_4 q^{n-1})} q^n \langle R_m, R_n \rangle_{\text{cher}} \delta_{m,n}.$$

**Proof.** We will use (2.5) and (2.11) to expand the  $T$ 's, so consider the integral

$$\begin{aligned} & \frac{1}{2\pi i} \oint_{|z|=1} (t_1 z, q t_1/z; q)_j (t_2 z, q t_2/z; q)_k w_{\text{cher}}(z; \mathbf{t} | q) \frac{dz}{z} \\ &= \frac{1}{2\pi i} \oint_{|z|=1} (t_1 z, t_1/z; q)_j (t_2 z, t_2/z; q)_k \frac{(z - q^k t_2)(1 - q^j t_1/z)}{z - 1/z} w_{\text{aw}}(z; \mathbf{t} | q) \frac{dz}{z} \\ &= \frac{(1 - t_1 t_2 q^{k+j})}{2\pi} \int_0^\pi w_{\text{aw}}(\cos \theta; t_1 q^j, t_2 q^k, t_3, t_4 | q) d\theta \\ &= \frac{(t_1 t_2 t_3 t_4 q^{k+j}; q)_\infty}{(q, t_1 t_2 q^{j+k+1}, t_1 t_3 q^j, t_1 t_4 q^j, t_2 t_3 q^k, t_2 t_4 q^k, t_3 t_4; q)_\infty} \end{aligned}$$

Using (2.5) to sum on  $j$  we find

$$\begin{aligned} & \frac{1}{2\pi i} \oint_{|z|=1} T_n(z; \mathbf{t} | q) (t_2 z, q t_2 / z; q)_k w_{\text{cher}}(z; \mathbf{t} | q) \frac{dz}{z} \\ &= \frac{(t_1 t_2 t_3 t_4 q^k; q)_\infty}{(t_1 t_2 q^{k+1}, t_1 t_3, t_1 t_4, t_2 t_3 q^k, t_2 t_4 q^k, t_3 t_4; q)_\infty} \\ & \quad \times {}_3\phi_2 \left( \begin{matrix} q^{-n}, t_1 t_2 t_3 t_4 q^{n-1}, t_1 t_2 q^{k+1} \\ q t_1 t_2, t_1 t_2 t_3 t_4 q^k \end{matrix} \middle| q, q \right) \end{aligned}$$

Again the  $q$ -Pfaff-Saalschütz theorem [7, (II.12)] evaluates the  ${}_3\phi_2$  as

$$\frac{(q^{-k}, q^{2-n}/t_3 t_4; q)_n}{(q t_1 t_2, q^{1-k-n}/t_1 t_2 t_3 t_4; q)_n}$$

which vanishes when  $k < n$ . This establishes orthogonality for  $m < n$ , and the  $k = n$  case establishes Proposition 4 for  $m = n$  using (2.11).  $\blacksquare$

To complete the proof of Theorem 1, we switch the orthogonality results for  $R_n$  and  $T_n$  to results for  $R_n$  and  $U_n$  using Proposition 1,

$$c_n U_n = T_n - R_n, \quad c_n = \frac{q t_1 (1 - q^{-n})(1 - t_1 t_2 t_3 t_4 q^{n-1})}{(1 - t_1 t_2)(1 - q t_1 t_2)(1 - t_1 t_3)(1 - t_1 t_4)}.$$

The only remaining orthogonality relation we must check is

$$\langle U_n, R_n \rangle_{\text{cher}} = c_n^{-1} \langle T_n - R_n, R_n \rangle_{\text{cher}} = c_n^{-1} (\langle T_n, R_n \rangle_{\text{cher}} - \langle R_n, R_n \rangle_{\text{cher}}) = 0.$$

The value of  $\langle U_n, U_n \rangle_{\text{cher}}$ , which is non-zero, is found using Proposition 4.

$$\begin{aligned} \langle U_n, U_n \rangle_{\text{cher}} &= c_n^{-2} \langle T_n - R_n, T_n - R_n \rangle_{\text{cher}} \\ &= c_n^{-2} (\langle T_n, T_n \rangle_{\text{cher}} - \langle R_n, R_n \rangle_{\text{cher}}) \\ &= c_n^{-2} \left( \frac{(1 - t_1 t_2)(1 - t_3 t_4 / q)}{(1 - t_1 t_2 q^n)(1 - t_3 t_4 q^{n-1})} q^n - 1 \right) \langle R_n, R_n \rangle_{\text{cher}} \\ &= c_n^{-2} \frac{(1 - q^n)(1 - t_1 t_2 t_3 t_4 q^{n-1})}{(1 - t_1 t_2 q^n)(1 - t_3 t_4 q^{n-1})} \langle R_n, R_n \rangle_{\text{cher}}. \end{aligned}$$

## 4 More orthogonal bases for $V_n$

In this section we find additional orthogonal bases for  $V_n$ , see Theorem 2, Theorem 3, and Theorem 4. We also explicitly give three and four term recurrence relations for some of these bases, which are the analogues of the recurrence relation for orthogonal polynomials.

The first result of this section is

**Theorem 2.** *The set  $\{T_0, \dots, T_n, S_1, \dots, S_n\}$  is an orthogonal basis for  $V_n$  with respect to the symmetric bilinear form  $\langle \cdot, \cdot \rangle_{\text{cher}}$ . Moreover,  $\langle T_m, T_m \rangle_{\text{cher}}$  and  $\langle S_m, S_m \rangle_{\text{cher}}$  are non-zero.*

**Proof.** Proposition 4 gives orthogonality of the  $T_n$ 's. By Proposition 1 there are linear dependencies amongst  $\{R_n, U_n, T_n\}$ , and  $\{R_n, U_n, S_n\}$ . So  $\langle S_m, S_m \rangle$ ,  $\langle T_m, T_m \rangle$ , and  $\langle S_m, T_m \rangle$  may all be changed to the appropriate linear combinations of  $R_n$ 's and  $T_m$ 's. The results are given in the next Proposition.  $\blacksquare$

**Proposition 5.** For all  $n \geq 1$  and  $m \geq 0$  we have the orthogonality relations

$$\begin{aligned} \langle S_n, T_m \rangle_{cher} &= 0, \\ t_1 \langle S_n, R_m \rangle_{cher} &= (1 - t_1 t_3)(1 - t_1 t_4) \langle R_n, R_m \rangle_{cher}. \end{aligned}$$

For all  $n \geq 1$  and  $m \geq 1$  we have the orthogonality relations

$$\langle S_n, S_m \rangle_{cher} = \frac{q^n (1 - t_1 t_3)^2 (1 - t_1 t_4)^2 (1 - t_1 t_2) (1 - t_3 t_4 / q)}{(q^n - 1) (1 - t_1 t_2 t_3 t_4 q^{n-1}) t_1^2} \langle R_n, R_m \rangle_{cher}.$$

**Proof.** This follows by direct calculation using Proposition 1, Proposition 3 and Proposition 4.  $\blacksquare$

**Corollary 1.** If  $n \geq 1$ , then for any  $f \in V_{n-1}$  we have

$$\langle R_n, f \rangle_{cher} = \langle S_n, f \rangle_{cher} = \langle T_n, f \rangle_{cher} = \langle U_n, f \rangle_{cher} = 0.$$

Using Corollary 1 one may explicitly give many orthogonal bases using the Askey-Wilson polynomials  $R_n$ . For example, for the basis

$$\{R_0, \dots, R_n, T_1, \dots, T_n\}$$

of  $V_n$ , the only pairwise non-orthogonalities involve  $T_m$  and  $R_m$ . One can replace these two polynomials by orthogonal linear combinations  $R_m + c_m T_m$  and  $R_m + d_m T_m$ . The values of the constants  $c_m$  and  $d_m$  yielding orthogonality can be chosen using Proposition 3 and Proposition 4. The non-symmetric Askey-Wilson polynomial of §6 are one way of accomplishing this.

Another such choice corresponds to applying the Gram-Schmidt process to the ordered basis  $\{1, z^{-1}, z^1, \dots, z^{-n}, z^n\}$  on  $V_n$  with  $\langle, \rangle_{cher}$ .

**Theorem 3.** An orthogonal basis for  $V_n$  is  $\{X_0, X_{-1}, X_1, \dots, X_{-n}, X_n\}$ , where

$$\begin{aligned} X_{-n} &= (1 - t_1 t_2)(1 - t_1 t_3)(1 - t_1 t_4) R_n - t_1 (1 - t_1 t_2 t_3 t_4 q^{n-1}) S_n, \quad n > 0, \\ X_n &= (1 - t_1 t_2)(1 - t_1 t_3)(1 - t_1 t_4) R_n + t_1^2 t_2 (1 - q^n) S_n, \quad n \geq 0. \end{aligned}$$

**Proof.** The choice of the coefficients for  $X_{-n}$  allows  $X_{-n} \in V_{n-1} \oplus sp\{z^{-n}\}$ , while the coefficients for  $X_n$  force  $\langle X_n, X_{-n} \rangle_{cher} = 0$  via Proposition 5. An explicit computation shows that all of the  $L^2$ -norms are non-zero.  $\blacksquare$

For the ordered basis  $\{1, z^1, z^{-1}, \dots, z^n, z^{-n}\}$  on  $V_n$  we have an analogous result.

**Theorem 4.** An orthogonal basis for  $V_n$  is  $\{Y_0, Y_{-1}, Y_1, \dots, Y_{-n}, Y_n\}$ , where

$$\begin{aligned} Y_{-n} &= q^{n-1} t_3 t_4 (1 - t_1 t_2)(1 - t_1 t_3)(1 - t_1 t_4) R_n - t_1 (1 - t_1 t_2 t_3 t_4 q^{n-1}) S_n, \quad n > 0, \\ Y_n &= q^n (1 - t_1 t_2)(1 - t_1 t_3)(1 - t_1 t_4) R_n + t_1 (1 - q^n) S_n, \quad n \geq 0. \end{aligned}$$

Orthogonal bases from Theorem 3 and Theorem 4 have a three term recurrence relation.

**Proposition 6.** For  $n \geq 1$ , there exist constants  $a_n, b_n$  and  $c_n$  such that

$$zX_{-n}(z) = a_n X_{-n}(z) + b_n X_n(z) + c_n X_{n-1}(z).$$

**Proof.** Because  $X_{-n}$  has no  $z^n$  term,  $zX_{-n}(z) \in V_n$  and has an expansion

$$zX_{-n}(z) = c_0 X_0(z) = \sum_{i=1}^n (c_i X_i(z) + c_{-i} X_{-i}(z)).$$

By orthogonality, we can find the coefficients

$$c_k = \frac{\langle zX_{-n}, X_k \rangle_{cher}}{\langle X_k, X_k \rangle_{cher}} = \frac{\langle X_{-n}, zX_k \rangle_{cher}}{\langle X_k, X_k \rangle_{cher}},$$

as long as  $\langle X_k, X_k \rangle_{cher} \neq 0$  (which is true here). The only terms to survive the orthogonality are  $X_n$ ,  $X_{-n}$ , and  $X_{n-1}$ . The term  $X_{1-n}$  does not survive because  $zX_{1-n} \in V_{n-1}$ :

$$\langle zX_{-n}, X_{1-n} \rangle_{cher} = \langle X_{-n}, zX_{1-n} \rangle_{cher} = 0.$$

■

We record two Propositions for these recurrences. The coefficients may be found by considering specific powers of  $z$  on each side, we do not give the details of the computation.

**Proposition 7.** *The constants in Proposition 6 are*

$$\begin{aligned} a_n &= \frac{(-t_1 - t_2 + t_1t_2(t_3 + t_4)q^{n-1})(1 - t_1t_2t_3t_4q^{2n-1})}{(-1 - t_1t_2 + t_1t_2t_3t_4q^{n-1} + t_1t_2q^n)(1 - t_1t_2t_3t_4q^{2n-2})} \\ b_n &= -\frac{(-t_1 - t_2 + t_1t_2(t_3 + t_4)q^{n-1})(1 - t_1t_2t_3t_4q^{n-1})(1 - t_3t_4q^{n-1})}{(-1 - t_1t_2 + t_1t_2t_3t_4q^{n-1} + t_1t_2q^n)(1 - t_1t_2t_3t_4q^{2n-2})} \\ c_n &= \frac{t_1(1 - t_3t_4q^{n-1})(1 - t_2t_4q^{n-1})(1 - t_2t_3q^{n-1})}{(1 - t_1t_2q^{n-1})(1 - t_1t_2t_3t_4q^{2n-2})}. \end{aligned}$$

**Proposition 8.** *For  $n \geq 1$  we have*

$$\begin{aligned} \frac{1}{z}X_n &= a_nX_{-n-1} + b_nX_n + c_nX_{-n}, \\ a_n &= \frac{(1 - t_1t_2q^n)(1 - t_1t_3q^n)(1 - t_1t_4q^n)}{t_1(1 - t_3t_4q^n)(1 - t_1t_2t_3t_4q^{2n})} \\ b_n &= \frac{(-t_1 - t_2 + t_1t_2(t_3 + t_4)q^n)(1 - t_1t_2t_3t_4q^{2n-1})}{(-1 - t_1t_2 + t_1t_2t_3t_4q^{n-1} + t_1t_2q^n)(1 - t_1t_2t_3t_4q^{2n})} \\ c_n &= -\frac{(1 - q^n)(1 - t_1t_2q^n)(-t_1 - t_2 + t_1t_2(t_3 + t_4)q^n)}{(-1 - t_1t_2 + t_1t_2t_3t_4q^{n-1} + t_1t_2q^n)(1 - t_1t_2t_3t_4q^{2n})}. \end{aligned}$$

There are four term relations for  $zX_n$  and  $\frac{1}{z}X_{-n}$ .

**Proposition 9.** *For  $n \geq 1$ , there exist constants  $a_n$ ,  $b_n$ ,  $c_n$  and  $d_n$  such that*

$$zX_n(z) = a_nY_{-n-1}(z) + b_nX_n(z) + c_nY_{-n}(z) + d_nX_{n-1}(z).$$

**Proof.** By orthogonality we have

$$zX_n(z) = a'_nX_{-n-1}(z) + b'_nX_{n+1}(z) + c'_nX_{-n}(z) + d'_nX_n(z) + e'_nX_{n-1}(z).$$

The sixth term  $X_{1-n}(z)$  does not appear because  $zX_{1-n} \in V_{n-1}$ :

$$\langle zX_n, X_{1-n} \rangle_{cher} = \langle X_n, zX_{1-n} \rangle_{cher} = 0.$$

Since  $sp\{X_{-n-1}, X_{n+1}\} = sp\{Y_{-n-1}, Y_{n+1}\}$ , the first two terms may be written as linear combination of  $Y_{-n-1}$  and  $Y_{n+1}$ , But  $zX_{-n}(z)$  has no  $z^{-n-1}$  term, thus only  $Y_{-n-1}$  appears. Because  $sp\{X_{-n}, X_n\} = sp\{Y_{-n}, X_n\}$  we can replace  $X_n$  by  $Y_{-n}$ . ■

**Proposition 10.** *The constants in Proposition 9 are*

$$\begin{aligned} a_n &= \frac{(1 - t_1 t_2 q^n)(1 - t_1 t_3 q^n)(1 - t_1 t_4 q^n)(-1 - t_1 t_2 + t_1 t_2 t_3 t_4 q^{n-1} + t_1 t_2 q^n)}{t_1(1 - t_3 t_4 q^n)(1 - t_1 t_2 t_3 t_4 q^{2n-1})(1 - t_1 t_2 t_3 t_4 q^{2n})}, \\ b_n &= \frac{q^n(-1 - t_1 t_2 + t_1 t_2 t_3 t_4 q^{n-1} + t_1 t_2 q^n)(-t_3 - t_4 + t_2 t_3 t_4 q^n + t_1 t_3 t_4 q^n)}{(1 - t_1 t_2 t_3 t_4 q^{2n-1})(1 - t_1 t_2 t_3 t_4 q^{2n})}, \\ c_n &= -\frac{(1 - q^n)(1 - t_1 t_2 q^n)(-t_1 - t_2 + t_1 t_2(t_3 + t_4)q^{n-1})}{(1 - t_1 t_2 t_3 t_4 q^{2n-2})(1 - t_1 t_2 t_3 t_4 q^{2n-1})}, \\ d_n &= \frac{t_1(1 - q^n)(1 - t_1 t_2 q^n)(1 - t_2 t_3 q^{n-1})(1 - t_2 t_4 q^{n-1})(1 - t_3 t_4 q^{n-1})}{(1 - t_1 t_2 q^{n-1})(1 - t_1 t_2 t_3 t_4 q^{2n-2})(1 - t_1 t_2 t_3 t_4 q^{2n-1})}. \end{aligned}$$

**Proposition 11.** *For  $n \geq 1$  we have*

$$\begin{aligned} \frac{1}{z}X_{-n} &= a_n X_{-n-1} + b_n X_{-n} + c_n Y_n + d_n Y_{n-1}, \\ a_n &= \frac{(1 - t_1 t_2 q^n)(1 - t_1 t_3 q^n)(1 - t_1 t_4 q^n)(1 - t_3 t_4 q^{n-1})(1 - t_1 t_2 t_3 t_4 q^{n-1})}{t_1(1 - t_3 t_4 q^n)(1 - t_1 t_2 t_3 t_4 q^{2n-1})(1 - t_1 t_2 t_3 t_4 q^{2n})}, \\ b_n &= \frac{q^{n-1}(-t_3 - t_4 + (t_1 + t_2)t_3 t_4 q^{n-1})(-1 - t_1 t_2 + t_1 t_2 t_3 t_4 q^{n-1} + t_1 t_2 q^n)}{(1 - t_1 t_2 t_3 t_4 q^{2n-2})(1 - t_1 t_2 t_3 t_4 q^{2n-1})}, \\ c_n &= -\frac{(1 - t_3 t_4 q^{n-1})(1 - t_1 t_2 t_3 t_4 q^{n-1})(-t_1 - t_2 + t_1 t_2(t_3 + t_4)q^n)}{(1 - t_1 t_2 t_3 t_4 q^{2n-1})(1 - t_1 t_2 t_3 t_4 q^{2n})}, \\ d_n &= \frac{t_1(1 - t_2 t_3 q^{n-1})(1 - t_2 t_4 q^{n-1})(1 - t_3 t_4 q^{n-1})(-1 - t_1 t_2 + t_1 t_2 t_3 t_4 q^{n-1} + t_1 t_2 q^n)}{(1 - t_1 t_2 q^{n-1})(1 - t_1 t_2 t_3 t_4 q^{2n-2})(1 - t_1 t_2 t_3 t_4 q^{2n-1})}. \end{aligned}$$

The two bases of Theorem 3 and Theorem 4 can be related, using (2.12). To explicitly show the parameter dependence we write  $X_n(z; \mathbf{t}|q)$  for  $X_n$  and  $Y_n(z; \mathbf{t}|q)$  for  $Y_n$ .

**Proposition 12.** *If  $n \geq 1$ , we have*

$$\begin{aligned} X_{-n}(1/z; 1/\mathbf{t}|1/q) &= -Y_{-n}(z; \mathbf{t}|q)/t_1^3 t_2^2 t_3^2 t_4^2 q^{n-1}, \\ X_n(1/z; 1/\mathbf{t}|1/q) &= -Y_n(z; \mathbf{t}|q)/t_1^3 t_2 t_3 t_4 q^n. \end{aligned}$$

Thus three term relations can be explicitly given for  $zY_n$  and  $\frac{1}{z}Y_{-n}$ , and four term relations for  $zY_{-n}$  and  $\frac{1}{z}Y_n$ .

## 5 The operators $A_0$ and $A_1$

Motivated by operators previously given in Noumi-Stokman [17], in this section we define linear transformations  $A_0$  and  $A_1$  on  $V_n$ . We explicitly find their actions on the possible bases  $R_m$ ,  $S_m$ ,  $T_m$ , and  $U_m$ . We use the self-adjointness of these operators to give an alternative proof of Theorem 1, Theorem 2, Theorem 3, and Theorem 4. We describe the eigenvalue problem whose solutions are the non-symmetric Askey-Wilson polynomials. We explicitly give the creation operators of Noumi-Stokman [17] via the three and four term recurrence relations of §4. These results are in [17], but we include them for completeness.

For a Laurent polynomial  $f \in V_n$ , define

$$(A_0 f)(z) = \frac{(1 - t_3/z)(1 - t_4/z)}{(1 - q/z^2)}(f(q/z) - f(z)),$$

and

$$(A_1 f)(z) = \frac{(1 - t_1 z)(1 - t_2 z)}{(1 - z^2)}(f(1/z) - f(z)).$$

By a direct computation it is easy to see that each  $A_i$  is a linear transformation from  $V_n$  to  $V_n$ .

**Proposition 13.** *Each  $A_i : V_n \rightarrow V_n$  is self-adjoint, that is for any  $f, g \in V_n$ , we have*

$$\langle A_0 f, g \rangle_{cher} = \langle f, A_0 g \rangle_{cher}, \quad \langle A_1 f, g \rangle_{cher} = \langle f, A_1 g \rangle_{cher} .$$

**Proof.** It is clear that  $\langle A_0 f, g \rangle_{cher}$  equals

$$\begin{aligned} & \frac{1}{2\pi i} \oint_{|z|=1} \frac{(q, z^2, 1/z^2; q)_\infty}{\prod_{j=1}^4 (t_j z, t_j/z; q)_\infty} \frac{(1-t_1/z)(1-t_2/z)}{1-z^{-2}} \\ & \quad \times \frac{(1-t_3/z)(1-t_4/z)}{(1-q/z^2)} [f(q/z)g(z) - f(z)g(z)] \frac{dz}{z} \\ &= \frac{1}{2\pi i} \oint_{|z|=1} \frac{(q, z^2, q^2/z^2; q)_\infty}{\prod_{j=1}^4 (t_j z, qt_j/z; q)_\infty} [f(q/z)g(z) - f(z)g(z)] \frac{dz}{z} \end{aligned}$$

This is a difference of two integrals. In the first we let  $z \rightarrow q/z$ . A calculation shows that the first integral becomes

$$\frac{1}{2\pi i} \oint_{|z|=q} \frac{(q, z^2, q^2/z^2; q)_\infty}{\prod_{j=1}^4 (t_j z, qt_j/z; q)_\infty} f(z)g(q/z) \frac{dz}{z} .$$

We now deform the above contour to the circular contour  $|z| = 1$  because the integrand has no poles between the contours since each  $|t_i| < 1$ . The proof of  $\langle A_1 f, g \rangle_{cher} = \langle f, A_1 g \rangle_{cher}$  is similar and will be omitted.  $\blacksquare$

Note that in the proof of Proposition 13 we only used the fact that  $f$  and  $g$  are analytic in  $q \leq |z| \leq 1/q$ .

**Proposition 14.** *The action of the operators  $A_0$  and  $A_1$  on the functions  $R_n, T_n, S_n$ , and  $U_n$  is given by*

$$\begin{aligned} A_0(R_n) &= \frac{t_1(q^{-n} - 1)(1 - t_1 t_2 t_3 t_4 q^{n-1})}{(1 - t_1 t_2)(1 - t_1 t_3)(1 - t_1 t_4)} S_n = \alpha_n S_n, & A_1(R_n) &= 0, \\ A_0(T_n) &= 0, & A_1(T_n) &= \frac{qt_1(q^{-n} - 1)(1 - t_1 t_2 t_3 t_4 q^{n-1})}{(1 - qt_1 t_2)(1 - t_1 t_3)(1 - t_1 t_4)} U_n = \beta_n U_n \\ A_0(S_n) &= (-1 + t_3 t_4/q) S_n, & A_1(S_n) &= \frac{q^{1-n}(1 - t_1 t_2 q^n)(1 - t_3 t_4 q^{n-1})}{(1 - qt_1 t_2)} U_n \\ A_0(U_n) &= q^{-1}(1 - qt_1 t_2) S_n, & A_1(U_n) &= (-1 + t_1 t_2) U_n, \end{aligned}$$

respectively.

**Proof.** These follow by direct term by term application of  $A_0$  and  $A_1$  to the defining series.  $\blacksquare$

These relations give another proof of the orthogonality relations in Theorem 1 and Theorem 2, for example

$$0 = \langle A_0(T_n), R_m \rangle_{cher} = \langle T_n, A_0(R_m) \rangle_{cher} = \alpha_m \langle T_n, S_m \rangle_{cher},$$

$$0 = \langle A_1(R_n), U_m \rangle_{cher} = \langle R_n, A_1(U_m) \rangle_{cher} = \beta_m \langle R_n, U_m \rangle_{cher} .$$

From Proposition 14 the operators  $A_0$  and  $A_1$  act on the 2-dimensional space  $W_n$  spanned by  $\{R_n, S_n, U_n, T_n\}$ . An elementary computation yields the next two Theorems.

**Theorem 5.** *If  $n \geq 1$ , the eigenvalues of  $(A_1 - t_1 t_2 I)(A_0 - q^{-1} t_3 t_4 I)$  on  $W_n$  are  $q^{-n}$  and  $t_1 t_2 t_3 t_4 q^{n-1}$ , with corresponding eigenvectors  $X_{-n}(z; \mathbf{t}|q)$  and  $Y_n(z; \mathbf{t}|q)$ .*

**Theorem 6.** *If  $n \geq 1$ , the eigenvalues of  $(A_0 - q^{-1} t_3 t_4 I)(A_1 - t_1 t_2 I)$  on  $W_n$  are  $q^{-n}$  and  $t_1 t_2 t_3 t_4 q^{n-1}$ , with corresponding eigenvectors  $Y_{-n}(z; \mathbf{t}|q)$  and  $X_n(z; \mathbf{t}|q)$ .*

These two Theorems give alternative proofs of Theorem 3 and Theorem 4. For example

$$\begin{aligned} q^{-n} \langle X_{-n}, X_n \rangle_{cher} &= \langle (A_1 - t_1 t_2 I)(A_0 - q^{-1} t_3 t_4 I) X_{-n}, X_n \rangle_{cher} \\ &= \langle X_{-n}, (A_0 - q^{-1} t_3 t_4 I)(A_1 - t_1 t_2 I) X_n \rangle_{cher} \\ &= t_1 t_2 t_3 t_4 q^{n-1} \langle X_{-n}, X_n \rangle_{cher} \end{aligned}$$

implying that  $0 = \langle X_{-n}, X_n \rangle_{cher}$ .

Our next goal is to define ladder maps  $\mathbb{S}_0$  and  $\mathbb{S}_1$  such that  $\mathbb{S}_1$  maps  $Y_n$  to  $X_{-n-1}$ , and  $\mathbb{S}_0$  maps  $X_{-n}$  to  $Y_n$ . If such maps are found, then  $Y_n$  is an  $n^{\text{th}}$  iterate of their composition  $\mathbb{S}_0 \mathbb{S}_1$  applied to  $Y_0$ , see Theorem 7 and Theorem 8.

Maps on  $W_n$  that interchange the bases elements  $\{X_{-n}, Y_n\}$  and  $\{X_n, Y_{-n}\}$  can be defined using the commutators of  $A_i$  with

$$Y = (A_1 - t_1 t_2 I)(A_0 - q^{-1} t_3 t_4 I) \text{ and } \mathbb{Y} = (A_0 - q^{-1} t_3 t_4 I)(A_1 - t_1 t_2 I).$$

From Proposition 14 the actions of  $Y, \mathbb{Y}, A_0$  and  $A_1$  are explicitly known on the 2-dimensional space  $W_n = \text{span}\{R_n, S_n, U_n, T_n\}$ . An explicit computation yields the next two Propositions.

**Proposition 15.** *For  $n \geq 1$  we have*

$$\begin{aligned} [Y, A_1](X_{-n}) &= t_1 t_2 q^{-n} (1 - t_3 t_4 q^{n-1}) (1 - t_1 t_2 t_3 t_4 q^{n-1}) Y_n, \\ [\mathbb{Y}, A_0](Y_{-n}) &= -q^{-1} t_3 t_4 (1 - t_3 t_4 q^{n-1}) (1 - t_1 t_2 t_3 t_4 q^{n-1}) X_n, \end{aligned}$$

and if  $n \geq 0$ ,

$$\begin{aligned} [Y, A_1](Y_n) &= q^{-n} (1 - q^n) (1 - t_1 t_2 q^n) X_{-n}, \\ [\mathbb{Y}, A_0](X_n) &= -q^{-2n} (1 - q^n) (1 - t_1 t_2 q^n) Y_{-n}. \end{aligned}$$

**Proposition 16.** *For  $n \geq 1$  we have*

$$\begin{aligned} [Y, A_1](R_n) &= \frac{t_1 q^{1-2n} (1 - q^n) (1 - t_3 t_4 q^{n-1}) (1 - t_1 t_2 q^n) (1 - t_1 t_2 t_3 t_4 q^{n-1})}{(1 - t_1 t_2) (1 - q t_1 t_2) / (1 - t_1 t_3) / (1 - t_1 t_4)} U_n \\ [Y, A_1](U_n) &= q^{-1} t_2 (1 - t_1 t_4) (1 - t_1 t_2) (1 - q t_1 t_2) (1 - t_1 t_3) R_n. \\ [Y, A_1](S_n) &= -q^{-n} t_2 (1 - q^n t_1 t_2) (1 - t_1 t_3) (1 - t_1 t_4) (1 - t_3 t_4 q^{n-1}) R_n + \\ &\quad q^{1-2n} \frac{(1 - q^n) (1 - t_1 t_2 q^n) (1 - t_3 t_4 q^{n-1}) (1 - t_1 t_2 t_3 t_4 q^{n-1})}{(1 - t_1 t_2) (1 - q t_1 t_2)} U_n \\ [Y, A_1](T_n) &= -q^{-n} t_1 t_2 (1 - t_1 t_2 t_3 t_4 q^{n-1}) R_n + \\ &\quad q^{1-2n} t_1 \frac{(1 - q^n) (1 - t_1 t_2 q^n) (1 - t_3 t_4 q^{n-1}) (1 - t_1 t_2 t_3 t_4 q^{n-1})}{(1 - t_1 t_2) (1 - q t_1 t_2) (1 - t_1 t_3) (1 - t_1 t_4)} U_n. \end{aligned}$$

We see from Proposition 15 that the commutator  $\mathbb{S}_0 = [Y, A_1]$  does map  $X_{-n}$  to  $Y_n$ . To define an appropriate  $\mathbb{S}_1$ , we need another pair of operators.

We define operators  $\hat{T}_1, \hat{T}_2 : V_n \rightarrow V_{n+1}$  by

$$(\hat{T}_1 f)(z) = \frac{(A_1 f)(z) + f(z)}{z}, \quad (\hat{T}_2 f)(z) = z((A_0 f)(z) + f(z)).$$

**Proposition 17.** *The action of  $\hat{T}_1$  on  $Y_n$  and  $X_{-n}$  is given by*

$$\begin{aligned} \hat{T}_1(Y_n) &= \frac{(1 - q^n t_1 t_2)(1 - q^n t_1 t_3)(1 - q^n t_1 t_4)}{t_1(1 - q^n t_3 t_4)(1 - q^{2n} t_1 t_2 t_3 t_4)} X_{-n-1} \\ &\quad + \frac{(t_1 + t_2 - q^n t_1 t_2(t_3 + t_4))}{(1 - q^{2n} t_1 t_2 t_3 t_4)} Y_n, \end{aligned}$$

if  $n \geq 0$  and

$$\begin{aligned} \hat{T}_1(X_{-n}) &= q^{n-1} \frac{t_1 t_2 (-t_4 + t_3 (-1 + q^{n-1}(t_1 + t_2)t_4))}{(-1 + q^{2(n-1)} t_1 t_2 t_3 t_4)} X_{-n} \\ &\quad + \frac{t_1^2 t_2 (1 - q^{n-1} t_2 t_3)(1 - q^{n-1} t_2 t_4)(1 - q^{n-1} t_3 t_4)}{(1 - q^{n-1} t_1 t_2)(-1 + q^{2(n-1)} t_1 t_2 t_3 t_4)} Y_{n-1}, \end{aligned}$$

if  $n \geq 1$ , respectively.

**Proof.** (Sketch) Note that  $\hat{T}_1 = \frac{1}{z}(A_1 + I)$ . The action of  $A_1$  on any element of  $W_n$  is given by Proposition 14. The action of division by  $z$  is given by Proposition 8, Proposition 11, and their  $Y$  versions. Thus  $\hat{T}_1(Y_n)$  and  $\hat{T}_1(X_{-n})$  may be given explicitly.  $\blacksquare$

From Theorem 5 we know that  $Y_n$  and  $X_{-n}$  are eigenfunctions of  $Y$ , and can obtain Proposition 18 from Proposition 17.

**Proposition 18.** *For  $n \geq 0$  we have*

$$\begin{aligned} [Y, \hat{T}_1](Y_n) &= - \frac{(1 - t_1 t_2 q^n)(1 - t_1 t_3 q^n)(1 - t_1 t_4 q^n)}{t_1 q^{n+1}(1 - t_3 t_4 q^n)} X_{-n-1}, \\ [Y, \hat{T}_1](X_{-n-1}) &= - t_1^2 t_2 \frac{(1 - t_2 t_3 q^n)(1 - t_2 t_4 q^n)(1 - t_3 t_4 q^n)}{q^{n+1}(1 - t_1 t_2 q^n)} Y_n. \end{aligned}$$

Finally we define  $\mathbb{S}_1 = [Y, \hat{T}_1]$ , which maps  $Y_n$  to  $X_{-n-1}$ . The next Theorem is Proposition 9.3 in [17]. Recall that  $Y_0 = (1 - t_1 t_2)(1 - t_1 t_3)(1 - t_1 t_4)$  is a constant.

**Theorem 7.** *We have*

$$c_n Y_n = (\mathbb{S}_0 \mathbb{S}_1)^n(Y_0), \quad d_n X_{-n-1} = \mathbb{S}_1(\mathbb{S}_0 \mathbb{S}_1)^n(Y_0).$$

where

$$\begin{aligned} c_n &= (-t_2)^n q^{-n(n+1)}(t_1 t_2, t_1 t_3, t_1 t_4, t_1 t_2 t_3 t_4; q)_n, \\ d_n &= -(-t_2)^n q^{-(n+1)^2}(t_1 t_2, t_1 t_3, t_1 t_4; q)_{n+1}(t_1 t_2 t_3 t_4; q)_n / t_1(1 - t_3 t_4 q^n). \end{aligned}$$

Completely analogous results hold for

$$\hat{T}_2, \quad \mathbb{S}_2 = [\mathbb{Y}, A_0], \quad \mathbb{S}_3 = [\mathbb{Y}, \hat{T}_2]$$

which we state below. Note that Proposition 15 shows that  $\mathbb{S}_2$  interchanges  $Y_{-n}$  and  $X_n$ .

**Proposition 19.** *For  $n \geq 0$  we have*

$$\begin{aligned} \hat{T}_2(X_n) &= \frac{t_3 + t_4 - t_3 t_4 q^n(t_1 + t_2)}{(1 - q^{2n} t_1 t_2 t_3 t_4)} X_n \\ &\quad + \frac{(1 - q^n t_1 t_2)(1 - q^n t_1 t_3)(1 - q^n t_1 t_4)}{t_1 q^n(1 - q^n t_3 t_4)(1 - q^{2n} t_1 t_2 t_3 t_4)} Y_{-n-1} \\ \hat{T}_2(Y_{-n-1}) &= t_1 t_3 t_4 q^n \frac{(1 - q^n t_2 t_3)(1 - q^n t_2 t_4)(1 - q^n t_3 t_4)}{(1 - q^n t_1 t_2)(1 - q^{2n} t_1 t_2 t_3 t_4)} X_{-n} \\ &\quad - q^n t_3 t_4 \frac{(t_2 + t_1 - t_1 t_2 q^n(t_3 + t_4))}{(1 - q^{2n} t_1 t_2 t_3 t_4)} Y_{-n-1}. \end{aligned}$$

The map  $\mathbb{S}_3 = [\mathbb{Y}, \hat{T}_2]$  interchanges  $X_n$  and  $Y_{-n-1}$ .

**Proposition 20.** *For  $n \geq 0$  we have*

$$\begin{aligned} [\mathbb{Y}, \hat{T}_2](X_n) &= \frac{(1 - t_1 t_2 q^n)(1 - t_1 t_3 q^n)(1 - t_1 t_4 q^n)}{t_1 q^{2n+1}(1 - t_3 t_4 q^n)} Y_{-n-1}, \\ [\mathbb{Y}, \hat{T}_2](Y_{-n-1}) &= q^{-1} t_1 t_3 t_4 \frac{(1 - t_2 t_3 q^n)(1 - t_2 t_4 q^n)(1 - t_3 t_4 q^n)}{(1 - t_1 t_2 q^n)} X_n. \end{aligned}$$

Recall that  $X_0 = (1 - t_1 t_2)(1 - t_1 t_3)(1 - t_1 t_4)$  is a constant.

**Theorem 8.** *We have*

$$e_n X_n = (\mathbb{S}_2 \mathbb{S}_3)^n(X_0), \quad f_n Y_{-n-1} = \mathbb{S}_3(\mathbb{S}_2 \mathbb{S}_3)^n(X_0),$$

where

$$\begin{aligned} e_n &= (-t_3 t_4 / t_1)^n q^{-n(n+1)} (t_1 t_2, t_1 t_3, t_1 t_4, t_1 t_2 t_3 t_4; q)_n, \\ f_n &= (-t_3 t_4 / t_1)^n q^{-n(n+1)} (t_1 t_2, t_1 t_3, t_1 t_4; q)_{n+1} (t_1 t_2 t_3 t_4; q)_n / t_1 q^{2n+1} (1 - t_3 t_4 q^n). \end{aligned}$$

Since any two elements from  $\{X_n, X_{-n}, Y_n, Y_{-n}\}$  form a basis for  $W_n$ , any element of  $W_n$  has several such formulas.

## 6 Non-symmetric Askey-Wilson polynomials

The non-symmetric Askey-Wilson polynomials may be defined as the eigenfunctions given in Theorem 5, namely

$$\{Y_0, Y_1, \dots, Y_n, \dots\} \cup \{X_{-1}, X_{-2}, \dots, X_{-n}, \dots\}.$$

This set of Laurent polynomials also satisfy an biorthogonality relation which follows from Theorem 3 and Theorem 4.

Define a new bilinear form on  $V_n$  by

$$\langle f, g \rangle_{\text{cher}'} = \frac{1}{2\pi i} \oint_{|z|=1} f(z) g(1/z) w_{\text{cher}}(z; \mathbf{t} | q) \frac{dz}{z}.$$

Put

$$P_m(z, \mathbf{t} | q) = \begin{cases} Y_m(z, \mathbf{t} | q) & \text{if } m \geq 0, \\ X_m(z, \mathbf{t} | q) & \text{if } m < 0, \end{cases}$$

and  $P'_m(z, \mathbf{t} | q) = P_m(z, 1/\mathbf{t} | 1/q)$ .

The biorthogonality relation with respect to this bilinear form [16], [17, §6.7] is

**Corollary 2.** *For  $m \neq n$ ,  $\langle P_m, P'_n \rangle_{\text{cher}'} = 0$ .*

**Proof.** We use Proposition 12 to change the bilinear form  $\langle, \rangle_{\text{cher}'}$  to  $\langle, \rangle_{\text{cher}}$ . For example, if  $m, n < 0$ ,  $m \neq n$ , then

$$\langle X_m, X'_n \rangle_{\text{cher}'} = - \langle X_m, Y_n \rangle_{\text{cher}} / t_1^3 t_2^2 t_3^2 t_4^2 q^{n-1} = 0,$$

while if  $m < 0$  and  $n \geq 0$

$$\langle X_m, Y'_n \rangle_{\text{cher}'} = - \langle X_m, X_n \rangle_{\text{cher}} t_1^3 t_2 t_3 t_4 q^n = 0,$$

The only term which requires checking is for  $n > 0$ ,

$$\langle Y_n, X'_{-n} \rangle_{cher'} = - \langle Y_n, Y_{-n} \rangle_{cher} / t_1^3 t_2 t_3^2 t_4^2 q^{n-1} = 0$$

and

$$\langle X_{-n}, Y'_n \rangle_{cher'} = - \langle X_{-n}, X_n \rangle_{cher} t_1^3 t_2 t_3^2 t_4^2 q^n = 0$$

by Theorem 3 and Theorem 4. ■

The  $L^2$ -norms may also be found.

**Corollary 3.** *We have for  $n > 0$ ,*

$$\begin{aligned} \langle Y_n, Y'_n \rangle_{cher'} &= -q^n \langle R_n, R_n \rangle_{cher} / t_1^3 t_2 t_3 t_4 \\ &\quad \times \frac{(1-t_1 t_2)(1-t_1 t_2 q^n)(1-t_1 t_3)^2(1-t_1 t_4)^2(1-t_1 t_2 t_3 t_4 q^{2n-1})}{(1-t_1 t_2 t_3 t_4 q^{n-1})}. \\ \langle X_{-n}, X'_{-n} \rangle_{cher'} &= -t_1^3 t_2 t_3^2 t_4^2 q^{1-n} \langle R_n, R_n \rangle_{cher} \\ &\quad \times \frac{(1-t_1 t_2)(1-t_3 t_4 q^{n-1})(1-t_1 t_3)^2(1-t_1 t_4)^2(1-t_1 t_2 t_3 t_4 q^{2n-1})}{(1-q^n)}. \end{aligned}$$

**Proof.** Using Proposition 12 we have

$$\langle Y_n, Y'_n \rangle_{cher'} = -q^n \langle Y_n, X_n \rangle_{cher} / t_1^3 t_2 t_3 t_4,$$

and  $\langle Y_n, X_n \rangle_{cher}$  may be found using Theorem 3, Theorem 4 and Proposition 5. The proof for  $\langle X_{-n}, X'_{-n} \rangle_{cher'}$  is similar. ■

Koornwinder and Bouzeffour [13] gave a positive definite orthogonality relation for the non-symmetric Askey-Wilson polynomials..

Macdonald [16, (6.6.8), p. 167] gives an expression for  $P_m(z; \mathbf{t}|q)$  as linear combination of  $R_m(z; \mathbf{t}|q)$  and  $T_m(z; \mathbf{t}|q)$ . This is equivalent to the Noumi-Stokman expression using Proposition 1.

Another [17, §6.8] biorthogonal pair of bases with respect to  $\langle, \rangle_{cher'}$  is given by the bases

$$\{R_0, \dots, R_n, U_1, \dots, U_n\} \text{ and } \{R'_0, \dots, R'_n, U'_1, \dots, U'_n\}.$$

The  $U_n(z; \mathbf{t}|q)$  are the anti-symmetric Askey-Wilson polynomials. However since

$$R_n(1/z; 1/\mathbf{t}|1/q) = R_n(z; \mathbf{t}|q), \quad U_n(1/z; 1/\mathbf{t}|1/q) = U_n(z; \mathbf{t}|q)/t_1 t_2,$$

this orthogonality is equivalent to Theorem 1.

## 7 Askey-Wilson orthogonality

One may ask if there are orthogonality relations for a basis of  $V_n$  using the Askey-Wilson weight. By symmetry, one could take

$$\{R_0, \dots, R_n, (z-1/z), (z-1/z)R_1, \dots, (z-1/z)R_{n-1}\}.$$

We include a partial result in this direction.

**Theorem 9.** *We have the orthogonality relation*

$$\begin{aligned} & \frac{1}{2\pi i} \oint_{|z|=1} R_m(z; \mathbf{t} | q) T_n(z; \mathbf{t} | q) w_{\text{aw}}(x; \mathbf{t} | q) \frac{dz}{z} \\ &= \frac{(-t_1^2)^n q^{\binom{n}{2}} (1+q^n) (t_1 t_2 t_3 t_4 q^{2n}, q)_\infty}{2 \prod_{1 \leq j < k \leq 4} (t_j t_k; q)_\infty} \prod_{2 \leq j < k \leq 4} (t_j t_k; q)_n \delta_{m,n}, \end{aligned}$$

for  $m \geq n$ .

**Proof.** Consider the integral

$$\frac{1}{2\pi i} \oint_{|z|=1} R_m(z; \mathbf{t} | q) (t_1 z, q t_1/z; q)_k w_{\text{aw}}(x; \mathbf{t} | q) \frac{dz}{z},$$

for  $0 \leq k \leq n$ . The integral vanishes when  $k = 0$  by (2.1), so we assume  $k > 0$ . Write  $(t_1 z, q t_1/z; q)_k$  as  $(1 - t_1 z)(1 - t_1 q^k/z)(q t_1 z, q t_1/z; q)_{k-1}$ . Therefore the above integral is

$$\begin{aligned} & \frac{1}{2\pi} \int_{-\pi}^{\pi} R_m(e^{i\theta}; \mathbf{t} | q) (q t_1 e^{i\theta}, q t_1 e^{-i\theta}; q)_{k-1} \\ & \times [1 - t_1(1 + q^k) \cos \theta + t_1^2 q^k] w_{\text{aw}}(x; \mathbf{t} | q) d\theta, \end{aligned}$$

which vanishes for  $k < m$  by (2.1). If  $k = m$  the above integral is

$$\begin{aligned} & \frac{1 + q^n}{4\pi} \int_{-\pi}^{\pi} R_m(e^{i\theta}; \mathbf{t} | q) (t_1 e^{i\theta}, t_1 e^{-i\theta}; q)_n w_{\text{aw}}(x; \mathbf{t} | q) d\theta \\ &= \frac{1 + q^n}{4\pi} \frac{(q, t_1 t_2, t_1, t_3, t_1 t_4; q)_n}{q^n (q^{-n}, t_1 t_2 t_3 t_4 q^{n-1}; q)_n} \int_{-\pi}^{\pi} R_m^2(e^{i\theta}) w_{\text{aw}}(x; \mathbf{t} | q) d\theta, \end{aligned}$$

and the result follows from (1.1), (2.1) and (2.3). ■

## 8 Recurrences

In this section we record three term recurrence relations satisfied by  $R_n, S_n, T_n$  and  $U_n$ . The three term recurrence relation for  $R_n$  is the Askey-Wilson recurrence relation [12, (3.1.4)]

$$\begin{aligned} [z + 1/z] R_n(z; \mathbf{t} | q) &= A_n^{(r)} R_{n+1}(z; \mathbf{t} | q) + C_n^{(r)} R_{n-1}(z; \mathbf{t} | q) \\ &+ [-A_n^{(r)} - C_n^{(r)} + t_1 + 1/t_1] R_n(z; \mathbf{t} | q), \end{aligned}$$

where

$$\begin{aligned} A_n^{(r)} &= \frac{(1 - t_1 t_2 t_3 t_4 q^{n-1}) \prod_{j=2}^4 (1 - t_1 t_j q^n)}{t_1 (1 - t_1 t_2 t_3 t_4 q^{2n-1}) (1 - t_1 t_2 t_3 t_4 q^{2n})}, \\ C_n^{(r)} &= \frac{t_1 (1 - q^n) \prod_{2 \leq j < k \leq 4} (1 - t_j t_k q^{n-1})}{(1 - t_1 t_2 t_3 t_4 q^{2n-2}) (1 - t_1 t_2 t_3 t_4 q^{2n-1})}. \end{aligned}$$

The three term recurrence relation for  $T_n$  is

$$\begin{aligned} [z + q/z] T_n(z; \mathbf{t} | q) &= A_n^{(t)} T_{n+1}(z; \mathbf{t} | q) + C_n^{(t)} T_{n-1}(z; \mathbf{t} | q) \\ &+ [-A_n^{(t)} - C_n^{(t)} + q t_1 + 1/t_1] T_n(z; \mathbf{t} | q), \end{aligned}$$

with  $n > 0$ , where

$$\begin{aligned} A_n^{(t)} &= \frac{(1 - t_1 t_2 t_3 t_4 q^{n-1}) (1 - t_1 t_2 q^{n+1}) \prod_{j=3}^4 (1 - t_1 t_j q^n)}{t_1 (1 - t_1 t_2 t_3 t_4 q^{2n-1}) (1 - t_1 t_2 t_3 t_4 q^{2n})}, \\ C_n^{(t)} &= \frac{q t_1 (1 - q^n) (1 - t_3 t_4 q^{n-2}) \prod_{j=3}^4 (1 - t_2 t_j q^{n-1})}{(1 - t_1 t_2 t_3 t_4 q^{2n-2}) (1 - t_1 t_2 t_3 t_4 q^{2n-1})}. \end{aligned}$$

On the other hand the three term recurrence relation for  $U_n$  is

$$[z + q/z]U_n(z; \mathbf{t}|q) = A_n^{(u)}U_{n+1}(z; \mathbf{t}|q) + C_n^{(u)}U_{n-1}(z; \mathbf{t}|q) \\ + [-A_n^{(u)} - C_n^{(u)} + qt_1 + q^{-1}/t_1]U_n(z; \mathbf{t}|q),$$

with  $n > 1$ , where

$$A_n^{(u)} = \frac{(1 - t_1t_2t_3t_4q^n)(1 - t_1t_2q^{n+1}) \prod_{j=3}^4 (1 - t_1t_jq^n)}{qt_1(1 - t_1t_2t_3t_4q^{2n-1})(1 - t_1t_2t_3t_4q^{2n})}, \\ C_n^{(u)} = \frac{qt_1(1 - q^{n-1})(1 - t_1t_2q^n)(1 - t_3t_4q^{n-2})}{(1 - t_1t_2t_3t_4q^{2n-2})(1 - t_1t_2t_3t_4q^{2n-1})} \\ \times \prod_{j=3}^4 (1 - t_1t_jq^{n-1})(1 - t_2t_jq^{n-1}).$$

Finally the three term recurrence relation for  $S_n$  is

$$[q^{-1/2}z + q^{1/2}/z]R_n(z; \mathbf{t}|q) = A_n^{(s)}R_{n+1}(z; \mathbf{t}|q) + C_n^{(s)}R_{n-1}(z; \mathbf{t}|q) \\ + [-A_n^{(s)} - C_n^{(s)} + t_1 + 1/t_1]R_n(z; \mathbf{t}|q),$$

and the coefficients are given by

$$A_n^{(s)} = \frac{(1 - t_1t_2t_3t_4q^n) \prod_{j=2}^4 (1 - t_1t_jq^n)}{t_1(1 - t_1t_2t_3t_4q^{2n-1})(1 - t_1t_2t_3t_4q^{2n})}, \\ C_n^{(s)} = \frac{t_1(1 - q^n) \prod_{2 \leq j < k \leq 4} (1 - t_jt_kq^{n-1})}{(1 - t_1t_2t_3t_4q^{2n-2})(1 - t_1t_2t_3t_4q^{2n-1})}.$$

## 9 Asymptotics

Consider a general balanced terminating  ${}_4\phi_3$ ,

$${}_4\phi_3 \left( \begin{matrix} q^{-n}, Aq^{n-1}, B, C \\ D, E, F \end{matrix} \middle| q, q \right), \quad \text{with } ABC = DEF.$$

Since the  ${}_4\phi_3$  is symmetric in  $B$  and  $C$  we may assume

$$|B| \leq |C|.$$

Ismail and Wilson [10] determined the large degree asymptotics of the Askey–Wilson polynomials. When  $|B| < |C|$  their result is

$${}_4\phi_3 \left( \begin{matrix} q^{-n}, Aq^{n-1}, B, C \\ D, E, F \end{matrix} \middle| q, q \right) = C^n \frac{(B, D/C, E/C, F/C; q)_\infty}{(B/C, D, E, F; q)_\infty} [1 + \mathcal{O}(q^{n/2})].$$

On the other hand when  $|B| = |C|$  we let  $B/C = e^{2i\theta}$  and in this case the Ismail–Wilson asymptotic result is

$${}_4\phi_3 \left( \begin{matrix} q^{-n}, Aq^{n-1}, B, C \\ D, E, F \end{matrix} \middle| q, q \right) = [1 + \mathcal{O}(q^{n/2})] \\ \times \left[ \frac{C^n (B, D/C, E/C, F/C; q)_\infty}{(B/C, D, E, F; q)_\infty} + \frac{B^n (C, D/B, E/B, F/B; q)_\infty}{(C/B, D, E, F; q)_\infty} \right].$$

We now record the asymptotics of  $R_n, S_n, T_n$  and  $U_n$ .

**Theorem 10.** For  $|z| = 1$  the following asymptotic formulas hold

$$\begin{aligned}
\prod_{j=2}^4 (t_1 t_j; q)_\infty R_n(z; \mathbf{t}) &= \left(\frac{t_1}{z}\right)^n \frac{\prod_{j=1}^4 (t_j z; q)_\infty}{(z^2; q)_\infty} [1 + \mathcal{O}(q^{n/2})] \\
&\quad + (t_1 z)^n \frac{\prod_{j=1}^4 (t_j/z; q)_\infty}{(1/z^2; q)_\infty} [1 + \mathcal{O}(q^{n/2})], \\
\frac{(qz^{-2}; q)_\infty \prod_{j=2}^4 (qt_1 t_j; q)_\infty}{(qt_1/z, qt_2/z, t_3/z, t_4/z; q)_\infty} S_n(z; \mathbf{t}) &= t_1^{n-1} z^n [1 + \mathcal{O}(q^{n/2})], \\
\frac{(qz^{-2}; q)_\infty (qt_1 t_2; q)_\infty \prod_{j=3}^4 (t_1 t_j; q)_\infty}{\prod_{j=1}^2 (qt_j/z, t_{j+2}/z; q)_\infty} T_n(z; \mathbf{t}) &= (t_1 z)^n [1 + \mathcal{O}(q^{n/2})], \\
\frac{\prod_{j=2}^4 (qt_1 t_j; q)_\infty}{(1 - qt_1 t_2)} U_n(z; \mathbf{t}) &= \frac{(qt_1)^{n-1} \prod_{j=1}^4 (t_j z; q)_\infty}{z^n (z^2; q)_\infty} [1 + \mathcal{O}(q^{n/2})] \\
&\quad + (qt_1)^{n-1} z^{n-2} \frac{(1 - t_1 z)(1 - t_2 z) \prod_{j=1}^4 (t_j/z; q)_\infty}{(1/z^2; q)_\infty (1 - t_1/z)(1 - t_2/z)} [1 + \mathcal{O}(q^{n/2})].
\end{aligned}$$

## 10 Discrete orthogonality

The Askey-Wilson polynomials have a discrete orthogonality, the  $q$ -Racah orthogonality [2], [12], when  $t_1 t_j = q^{-N}$ , for some  $j = 2, 3$  or  $4$ , and for some positive integer  $N$ . In this section we give the corresponding discrete orthogonality for the Laurent polynomials. This may be done by deforming the contours, we do not give the details.

The  $q$ -Racah measure for the Askey-Wilson polynomials is purely discrete and based upon the very well poised terminating  ${}_6\phi_5$  evaluation [7, (II.20)]

$$\begin{aligned}
\sum_{k=0}^N \frac{(t_1^2, t_1 t_2, t_1 t_3, t_1 t_4; q)_k}{(q, qt_1/t_2, qt_1/t_3, qt_1/t_4; q)_k} \frac{1 - t_1^2 q^{2k}}{1 - t_1^2} \left(\frac{q}{t_1 t_2 t_3 t_4}\right)^k \\
= \frac{(qt_1^2, qt_1^2/t_2 t_3, qt_1^2/t_2 t_4, qt_1^2/t_3 t_4; q)_\infty}{(qt_1/t_2, qt_1/t_3, qt_1/t_4, q/t_1 t_2 t_3 t_4; q)_\infty}.
\end{aligned} \tag{10.1}$$

Using

$$\frac{1 - t_1^2 q^{2k}}{1 - t_1^2} = \frac{1}{(1 - t_1^2)(1 - t_1 t_2)} (-t_1 t_2 (1 - q^k)(1 - q^k t_1/t_2) + (1 - t_1^2 q^k)(1 - t_1 t_2 q^k))$$

one can rewrite the left side of the  ${}_6\phi_5$  evaluation (10.1) as a sum of two  ${}_4\phi_3$ 's

$$\sum_{k=1}^N \frac{(qt_1^2, qt_1 t_2; q)_{k-1} (t_1 t_3, t_1 t_4; q)_k}{(qt_1/t_2, q; q)_{k-1} (qt_1/t_3, qt_1/t_4; q)_k} \left(\frac{q}{t_1 t_2 t_3 t_4}\right)^k (-t_1 t_2) +$$

$$\sum_{k=0}^N \frac{(qt_1^2, qt_1 t_2, t_1 t_3, t_1 t_4; q)_k}{(qt_1/t_2, qt_1/t_3, qt_1/t_4, q; q)_k} \left(\frac{q}{t_1 t_2 t_3 t_4}\right)^k.$$

**Definition 1.** If  $t_1 t_j = q^{-N}$  for some positive integer  $N$ , the Racah bilinear form on  $V_N$  is

defined by

$$\begin{aligned} \langle f, g \rangle_{\text{Racah}} = & \\ & \sum_{k=1}^N \frac{(qt_1^2, qt_1t_2; q)_{k-1}(t_1t_3, t_1t_4; q)_k}{(qt_1/t_2, q; q)_{k-1}(qt_1/t_3, qt_1/t_4; q)_k} \left( \frac{q}{t_1t_2t_3t_4} \right)^k (-t_1t_2)f(t_1q^k)g(t_1q^k) + \\ & \sum_{k=0}^N \frac{(qt_1^2, qt_1t_2, t_1t_3, t_1t_4; q)_k}{(qt_1/t_2, qt_1/t_3, qt_1/t_4, q; q)_k} \left( \frac{q}{t_1t_2t_3t_4} \right)^k f(q^{-k}/t_1)g(q^{-k}/t_1). \end{aligned}$$

**Theorem 11.** *If  $t_1t_j = q^{-N}$  for some positive integer  $N$  and  $j = 3$  or  $4$ , then*

$$\begin{aligned} & \{R_0, \dots, R_N, U_1, \dots, U_N\}, \{T_0, \dots, T_N, S_1, \dots, S_N\} \\ & \{X_{-N}, \dots, X_0, X_1, \dots, X_N\} \text{ and } \{Y_{-N}, \dots, Y_0, Y_1, \dots, Y_N\} \end{aligned}$$

are Racah-orthogonal bases for  $V_N$ .

If  $t_1t_2 = q^{-N}$ , the Racah bilinear form contains a zero term: the  $k = N$  term in the second sum is 0 due to  $(qt_1t_2; q)_k$ . In this case  $U_N$  and  $T_N$  are not well-defined. A slight modification of Theorem 11 can be formulated, we do not give it here.

## Acknowledgements

The research of Mourad E. H. Ismail is partially supported by Research Grants Council of Hong Kong under contracts #101410 and #101411 and by King Saud University, Riyadh through grant DSFP/MATH 01.

## References

- [1] G. E. Andrews, R. Askey and R. Roy, *Special Functions*, Cambridge University Press, Cambridge, 1999.
- [2] R. Askey and J. Wilson, A set of orthogonal polynomials that generalize the Racah coefficients or  $6 - j$  symbols. *SIAM J. Math. Anal.* **10** (1979), 10081016.
- [3] R. Askey and J. Wilson, Some basic hypergeometric orthogonal polynomials that generalize Jacobi polynomials, *Memoirs Amer. Math. Soc.* Number 317, 1985.
- [4] C. Brezinski, *Biorthogonality and its Applications to Numerical Analysis*, Monographs and Textbooks in Pure and Applied Mathematics, 156. Marcel Dekker, Inc., New York, 1992.
- [5] A. Bultheel, P. Gonzalez-Vera, E. Hendriksen, O. Njstad, Orthogonal rational functions and continued fractions, in *Special functions 2000: current Perspective and Future Directions*, NATO Sci. Ser. II Math. Phys. Chem., 30, Kluwer Acad. Publ., Dordrecht, 2001, pp. 87109.
- [6] I. Cherednik, Nonsymmetric Macdonald polynomials, *Internat. Math. Res. Notices* (1995), no. 10, 483-515.
- [7] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, second edition, Cambridge University Press, Cambridge, 2004.
- [8] M. E. H. Ismail, *Classical and Quantum Orthogonal Polynomials in One Variable*, Cambridge University Press, Cambridge, 2005.
- [9] M. E. H. Ismail and D. Masson, Generalized orthogonality and continued fractions, *J. Approx. Theory* **83** (1995), 1-40.
- [10] M. E. H. Ismail and J. A. Wilson, Asymptotic and generating relations for the  $q$ -Jacobi and  ${}_4\phi_3$  polynomials, *J. Approx. Theory* **36**, 1982, 43-54.
- [11] W. B. Jones, and W. J. Thron, *Continued Fractions. Analytic theory and applications*, Addison-Wesley Publishing Co., Reading, Mass., 1980, reissued by Cambridge University Press, Cambridge.
- [12] R. Koekoek and R. Swarttouw, The Askey-scheme of hypergeometric orthogonal polynomials and its  $q$ -analogues, *Reports of the Faculty of Technical Mathematics and Informatics* no. 98-17, Delft University of Technology, Delft, 1998.

- 
- [13] T. Koornwinder and F. Bouzeffour, Non-symmetric Askey-Wilson polynomials as vector-valued polynomials, *Applicable Analysis* **90** (2011), 731-746.
  - [14] L. Lorentzen and H. Waadeland, *Continued Fractions with Applications*, North-Holland Publishing Co., Amsterdam, 1992.
  - [15] L. Lorentzen and H. Waadeland, *Continued Fractions*, Vol. 1. Convergence theory. Second edition. Atlantis Press, Paris; World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2008.
  - [16] I. G. Macdonald, *Affine Hecke Algebras and Orthogonal Polynomials*, Cambridge University Press, Cambridge, 2003.
  - [17] M. Noumi and J. Stokman, Askey-Wilson polynomials: an affine Hecke algebra approach. Laredo Lectures on Orthogonal Polynomials and Special Functions, 111–144, *Adv. Theory Spec. Funct. Orthogonal Polynomials*, Nova Sci. Publ., Hauppauge, NY, 2004.
  - [18] V. Spiridonov and A. Zhedanov, Spectral transformation chains and some new biorthogonal rational functions, *Comm. Math. Phys.* **210** (2000), 49–83.
  - [19] H. Wilkinson, *The Algebraic Eigenvalue Problem*, Oxford University Press, Oxford, 1965, reprinted 1999.
  - [20] J. A. Wilson, private communication, 1980.
  - [21] A. Zhedanov, Biorthogonal rational functions and the generalized eigenvalue problem, *J. Approx. Theory* **101** (1999), 303–329.