

**MATH 5615H: ANALYSIS**  
**A SOLUTION TO PROBLEM 1 ON HW 6**

INSTRUCTOR: SASHA VORONOV

Here is a solution of Problem 1 on Homework 6.

**Solution:** Write down the statement that 0 is not a limit of the sequence  $\{na_n\}$ :

(1) there is an  $\epsilon > 0$  : for each  $N \geq 1$  there is an  $n \geq N$  :  $na_n \geq \epsilon$ .

We want to find a contradiction with this assumption. Of course, the contradiction will come from the fact that the series converges,

Let us get back to the statement about 0 not being a limit. We can find  $n_1$  such that  $n_1 a_{n_1} \geq \epsilon$ , then choose  $n_2 \geq n_1 + 1$  such that  $n_2 a_{n_2} \geq \epsilon$ , then choose  $n_3 \geq n_2 + 1$  such that  $n_3 a_{n_3} \geq \epsilon$ , and so on. This way we will get a subsequence of  $\{a_{n_k}\}$  such that  $n_k a_{n_k} \geq \epsilon$ . Then the  $n$ th partial sum  $s_{n_k}$  of the series  $\sum a_n$  can be estimated as follows:

$$\begin{aligned} s_{n_k} &= a_1 + \cdots + a_{n_1} + a_{n_1+1} + \cdots + a_{n_2} + \cdots + a_{n_k} \\ &\geq n_1 a_{n_1} + (n_2 - n_1) a_{n_2} + \cdots + (n_k - n_{k-1}) a_{n_k} \\ &\geq \epsilon \left( \frac{n_1}{n_1} + \frac{n_2 - n_1}{n_2} + \cdots + \frac{n_k - n_{k-1}}{n_k} \right). \end{aligned}$$

Here in the first inequality we used the fact that  $a_n \geq a_{n+1}$  for all  $n$ , and the second inequality came from the assumption about 0 not being a limit of  $\{a_n\}$ . We need to estimate the result so as to see that the series diverges.

**Digression:** One estimate can be done like that: since  $n_1 < n_2 < \cdots < n_k$ , we have

$$s_{n_k} \geq \epsilon \left( \frac{n_1}{n_k} + \frac{n_2 - n_1}{n_k} + \cdots + \frac{n_k - n_{k-1}}{n_k} \right) = \epsilon \frac{n_k}{n_k} = \epsilon.$$

Unfortunately, this does not contradict the convergence of the series. Therefore, we need to look for a finer estimate.

Let us continue with

(2)

$$s_{n_k} \geq \epsilon \left( \frac{n_1}{n_1} + \frac{n_2 - n_1}{n_2} + \cdots + \frac{n_k - n_{k-1}}{n_k} \right) = \epsilon \left( 1 + \left( 1 - \frac{n_1}{n_2} \right) + \cdots + \left( 1 - \frac{n_{k-1}}{n_k} \right) \right).$$

---

*Date:* October 31, 2020.

**Digression:** If we had  $(1+ \text{something})$  instead of  $(1- \text{something})$ , we would have  $s_{n_k} \geq \epsilon k$ , which would grow unboundedly. Unfortunately with the minuses, it does not work. We need to fine tune the estimate for  $s_{n_k}$ .

Note that we can use the fact that in (1) for any  $N \geq 1$ , there exists an  $n$ , in a better way. Namely, each time choosing  $n_{i+1}$  using (1), we can make sure it is at least  $2n_i$ .

**Digression: How can you arrive at this choice?** You stare at the estimate (2) and try to see, if there might be a reason why the last sum grows unboundedly. There is no immediate reason: for instance, all the  $n_{i-1}$ 's can be as close to their respective  $n_i$ 's as being adjacent, in which case  $n_{i-1}/n_i$  will be close to 1, and the  $i$ th term in the sum will be close to  $1/n_i$ , *i.e.*, too small. However, in this case, that is to say, if the  $n_i$ 's happened to be successive naturals, the partial sum will be a partial sum of the harmonic series  $\sum 1/n$ , which also diverges, but then there is a question about how to deal with intermediate situations, when the  $n_i$ 's are not too close to each other and not too far. Thus, it is better to try to see if fiddling with the choice of  $n_i$ 's, one can make each term in (2) large enough. How large? No way to make them larger than 1. On the other hand, 0 is not large enough. What is the most natural choice then? One half.

So, choosing  $n_{i+1} \geq 2n_i$  as per (1), we get

$$1 - \frac{n_i}{n_{i+1}} \geq 1 - \frac{n_i}{2n_i} = \frac{1}{2},$$

and

$$s_{n_k} > \frac{\epsilon k}{2}.$$

This implies that the sequence of partial sums diverges and so does the series.